Journal of Convex Analysis
Volume 21 (2014), No. 4, 989–1006

A Best Proximity Point Theorem for Generalized Weakly Contractive Non Self Mappings

A. Almeida
Departamento de Matemáticas, Universidad de Las Palmas de Gran Canaria, Campus de Tafira Baja, 35017 Las Palmas de Gran Canaria, Spain
aalmeida@dma.ulpgc.es

J. Harjani
Departamento de Matemáticas, Universidad de Las Palmas de Gran Canaria, Campus de Tafira Baja, 35017 Las Palmas de Gran Canaria, Spain
jharjani@dma.ulpgc.es

K. Sadarangani
Departamento de Matemáticas, Universidad de Las Palmas de Gran Canaria, Campus de Tafira Baja, 35017 Las Palmas de Gran Canaria, Spain
ksadaran@dma.ulpgc.es

Received: April 04, 2013
Revised manuscript received: July 16, 2013
Accepted: September 20, 2013

The purpose of this paper is to provide sufficient conditions for the existence of a unique best proximity point for generalized weakly contractive mappings in the context of complete metric spaces.

Keywords: Best proximity point, generalized weakly contractive mapping, fixed point

1991 Mathematics Subject Classification: 41A65, 47H10

1. Introduction

Let A and B nonempty subsets of a metric space (X, d). An operator $T\colon A \to B$ is said to be a k-contraction if there exists a $k \in [0, 1)$ such that $d(Tx, Ty) \leq kd(x, y)$ for any $x, y \in A$. Banach's contraction principle states that, when A is a complete subset of X and T is a k-contraction which maps A into itself then T has a unique fixed point in A.

Due to its wide applications in various fields, a huge number of generalizations and extended versions of this principle appear in the literature (see [8, 9]).

The following notion of a weakly contractive mapping was introduced by Alber and Guerre-Delabriere in the context of Hilbert spaces in [2] and Rhoades proved that the result of [2] is also valid in complete metric spaces.

ISSN 0944-6532 / $ 2.50 © Heldermann Verlag

Definition 1.1 ([10]). Let (X, d) be a metric space and $T\colon X \to X$ a mapping. T is said to be weakly contractive if

$$d(Tx, Ty) \leq d(x, y) - \phi\left(d(x, y)\right), \quad \text{for all } x, y \in X,$$

where $\phi\colon [0, \infty) \to [0, \infty)$ is a continuous and nondecreasing function such that $\phi(t) = 0$ if and only if $t = 0$.

Notice that if $\phi(t) = kt$ with $k \in [0, 1)$ then T is a $(1 - k)$-contraction.

The following result is due to Rhoades [10].

Theorem 1.2 ([10]). *Let (X, d) be a complete metric space and $T\colon X \to X$ a weakly contractive mapping. Then T has a unique fixed point in X.*

In [4], Dutta and Choudhury present the following generalization of Theorem 1.2.

Theorem 1.3 ([4]). *Let (X, d) be a complete metric space and $T\colon X \to X$ a mapping satisfying*

$$\varphi\left(d(Tx, Ty)\right) \leq \varphi\left(d(x, y)\right) - \phi\left(d(x, y)\right), \quad \textit{for all } x, y \in X,$$

where $\varphi, \phi\colon [0, \infty) \to [0, \infty)$ are continuous and nondecreasing functions such that $\phi(t) = \varphi(t) = 0$ if and only if $t = 0$. Then T has a unique fixed point.

In this paper, we present a best proximity point theorem for generalized weakly contractive non self mappings.

First, we present a brief discussion about best proximity points

Let A be a nonempty subset of a metric space (X, d) and $T\colon A \to X$ a mapping. The solutions of the equation $Tx = x$ are fixed points of T. Consequently, $T(A) \cap A \neq \emptyset$ is a necessary condition for the existence of a fixed point for the operator T. If this necessary condition does not hold then $d(Tx, x) > 0$ for any $x \in A$ and the mapping $T\colon A \to X$ does not have any fixed point. In this setting, our aim is to find an element $x \in A$ such that $d(x, Tx)$ is a minimum. Best approximation theory has been developed in this direction.

In our context we consider two nonempty subsets A and B of a complete metric space and a mapping $T\colon A \to B$. A natural question is whether one can find an element $x_0 \in A$ such that $d(x_0, Tx_0) = \min\{d(x, Tx)\colon x \in A.\}$. Since $d(x, Tx) \geq d(A, B)$ for any $x \in A$, the optimal solution to this problem will be the one for which the value $d(A, B)$ is attained by the real valued function $\varphi\colon A \to \mathbb{R}$ given by $\varphi(x) = d(x, Tx)$.

Some results about best proximity points can be found in [1, 5, 7, 11, 12].

2. Preliminaries

Let A and B be two nonempty subsets of a metric space (X, d). Denote by A_0 and B_0 the following sets:

$$A_0 = \{x \in A\colon d(x, y) = d(A, B) \text{ for some } y \in B\},$$
$$B_0 = \{y \in B\colon d(x, y) = d(A, B) \text{ for some } x \in A\},$$

where $d(A, B) = \inf\{d(x, y)\colon x \in A \text{ and } y \in B\}$.

In [7], the authors present sufficient conditions which guarantee when the sets A_0 and B_0 are nonempty. Moreover, in [11] the authors proved that A_0 is contained in the boundary of A.

Now we present the notion of a generalized weakly contractive non self mapping.

Definition 2.1. Let A and B be two nonempty subsets of a metric space (X, d). A mapping $T\colon A \to B$ is said to be generalized weakly contractive if

$$\psi(d(Tx, Ty)) \leq \psi(M(x, y) - d(A, B)) - \phi(d(x, y)), \quad \text{for any } x, y \in A,$$

where $\psi\colon [0, \infty) \to [0, \infty)$ is a continuous and nondecreasing function such that $\psi(t) = 0$ if and only if $t = 0$ and $\phi\colon [0, \infty) \to [0, \infty)$ is a nondecreasing function such that $\phi(t) = 0$ if and only if $t = 0$, and where

$$M(x, y) := \max\left\{d(x, y), d(x, Tx), d(y, Ty), \frac{1}{2}[d(x, Ty) + d(y, Tx)]\right\}.$$

Notice that in the case $A = B$ every operator $T\colon A \to A$ satisfying the contractive condition appearing in Theorem 1.3 is a generalized weakly contractive mapping, since in this case $d(A, A) = 0$ and

$$\varphi(d(Tx, Ty)) \leq \varphi(d(x, y)) - \phi(d(x, y)) \leq \varphi(M(x, y)) - \phi(d(x, y)).$$

In [13], the author introduced the following definition.

Definition 2.2 ([13]). Let (A, B) be a pair of nonempty subsets of a metric space (X, d) with $A_0 \neq \emptyset$. Then the pair (A, B) is said to have the P-property if and only if for any $x_1, x_2 \in A_0$ and $y_1, y_2 \in B_0$

$$\left.\begin{array}{l} d(x_1, y_1) = d(A, B) \\ d(x_2, y_2) = d(A, B) \end{array}\right\} \Rightarrow d(x_1, x_2) = d(y_1, y_2).$$

It is easily seen that for any nonempty subset A of (X, d), the pair (A, A) has the P-property.

In [13], the author proves that any pair (A, B) of nonempty closed convex subsets of a real Hilbert space H satisfies the P-property.

3. Main results

We start this section presenting our main result.

Theorem 3.1. *Let (A, B) be a pair of nonempty closed subsets of a complete metric space (X, d) such that $A_0 \neq \emptyset$. Let $T\colon A \to B$ be a continuous generalized weakly contractive mapping satisfying $T(A_0) \subset B_0$. Suppose that the pair (A, B) has the P-property. Then there exists a unique $x^* \in A$ such that $d(x^*, Tx^*) = d(A, B)$.*

Proof. Since A_0 es nonempty, we take $x_0 \in A$. As $Tx_0 \in T(A_0) \subset B_0$, we can find $x_1 \in A_0$ such that $d(x_1, Tx_0) = d(A, B)$. Similarly, since $Tx_1 \in T(A_0) \subset B_0$, there exists $x_2 \in A_0$ such that $d(x_2, Tx_1) = d(A, B)$. Repeating this process, we obtain a sequence (x_n) in A_0 satisfying

$$d(x_{n+1}, Tx_n) = d(A, B), \text{ for any } n \in \mathbb{N}. \tag{1}$$

Since (A, B) has the P-property, we have that

$$d(x_{n+1}, x_n) = d(Tx_n, Tx_{n-1}), \text{ for any } n \in \mathbb{N}.$$

Now, suppose that there exists an $n_0 \in \mathbb{N}$ with $d(x_{n_0+1}, x_{n_0}) = 0$. In this case we have

$$0 = d(x_{n_0+1}, x_{n_0}) = d(Tx_{n_0}, Tx_{n_0-1}),$$

and, consequently, $Tx_{n_0-1} = Tx_{n_0}$. Therefore,

$$d(A, B) = d(x_{n_0}, Tx_{n_0-1}) = d(x_{n_0}, Tx_{n_0}),$$

and the part of existence is proved.

On the contrary, suppose that $d(x_{n+1}, x_n) > 0$ for any $n \in \mathbb{N}$. Taking into account the contractive condition and (1) we have

$$\begin{aligned} &\psi\left(d(x_{n+1}, x_n)\right) \\ &= \psi\left(d(Tx_n, Tx_{n-1})\right) \leq \psi\left(M(x_n, x_{n-1}) - d(A, B)\right) - \phi(d(x_n, x_{n-1})). \end{aligned} \tag{2}$$

We can distinguish four cases.

Case 1: Suppose that $M(x_n, x_{n-1}) = d(x_n, x_{n-1})$. In this case, from the last inequality and the fact that ψ is nondecreasing, we get

$$\begin{aligned} \psi\left(d(x_{n+1}, x_n)\right) &= \psi\left(d(Tx_n, Tx_{n-1})\right) \\ &\leq \psi\left(d(x_n, x_{n-1}) - d(A, B)\right) - \phi\left(d(x_n, x_{n-1})\right) \\ &\leq \psi\left(d(x_n, x_{n-1})\right) - \phi\left(d(x_n, x_{n-1})\right) \\ &\leq \psi\left(d(x_n, x_{n-1})\right), \end{aligned} \tag{3}$$

and, again, as ψ is nondecreasing,

$$d(x_{n+1}, x_n) \leq d(x_n, x_{n-1}).$$

Case 2: Suppose that $M(x_n, x_{n-1}) = d(x_n, Tx_n)$. Since $d(x_{n+1}, Tx_n) = d(A, B)$, we have

$$\begin{aligned} M(x_n, x_{n-1}) - d(A, B) &= d(x_n, Tx_n) - d(A, B) \\ &\leq d(x_n, x_{n+1}) + d(x_{n+1}, Tx_n) - d(A, B) \\ &= d(x_n, x_{n+1}). \end{aligned}$$

This and (2) give us

$$\begin{aligned} \psi\left(d(x_{n+1}, x_n)\right) &= \psi\left(d(Tx_n, Tx_{n-1})\right) \\ &\leq \psi\left(d(x_n, Tx_n) - d(A, B)\right) - \phi\left(d(x_n, x_{n-1})\right) \\ &\leq \psi\left(d(x_n, x_{n+1})\right) - \phi\left(d(x_n, x_{n-1})\right). \end{aligned}$$

Since $d(x_n, x_{n-1}) > 0$ and $\phi(t) = 0$ if and only if $t = 0$, the last inequality implies that

$$\psi\left(d(x_{n+1}, x_n)\right) \leq \psi\left(d(x_n, x_{n+1})\right) - \phi\left(d(x_n, x_{n-1})\right) < \psi\left(d(x_n, x_{n+1})\right),$$

which is a contradiction.

Case 3: Suppose that $M(x_n, x_{n-1}) = d(x_{n-1}, Tx_{n-1})$. In this case, by an argument similar to the one used in Case 2, we have

$$\begin{aligned} M(x_n, x_{n-1}) - d(A, B) &= d(x_{n-1}, Tx_{n-1}) - d(A, B) \\ &\leq d(x_{n-1}, x_n) + d(x_n, Tx_{n-1}) - d(A, B) = d(x_{n-1}, x_n), \end{aligned}$$

and, therefore,

$$\begin{aligned} \psi\left(d(x_{n+1}, x_n)\right) &= \psi\left(d(Tx_n, Tx_{n-1})\right) \\ &\leq \psi\left(d(x_{n-1}, Tx_{n-1}) - d(A, B)\right) - \phi\left(d(x_n, x_{n-1})\right) \\ &\leq \psi\left(d(x_{n-1}, x_n)\right) - \phi\left(d(x_n, x_{n-1})\right) \\ &< \psi\left(d(x_{n-1}, x_n)\right). \end{aligned} \tag{4}$$

By using the fact that ψ is nondecreasing,

$$d(x_{n+1}, x_n) < d(x_{n-1}, x_n).$$

Case 4: Suppose that $M(x_n, x_{n-1}) = \frac{1}{2}\left[d(x_n, Tx_{n-1}) + d(x_{n-1}, Tx_n)\right]$. In this case, we have

$$\begin{aligned} M(x_n, x_{n-1}) - d(A, B) &= \frac{1}{2}\left[d(x_n, Tx_{n-1}) + d(x_{n-1}, Tx_n)\right] - d(A, B) \\ &\leq \frac{1}{2}\left[d(A, B) + d(x_{n-1}, x_{n+1}) + d(x_{n+1}, Tx_n)\right] - d(A, B) \\ &= \frac{1}{2}\left[d(A, B) + d(x_{n-1}, x_{n+1}) + d(A, B)\right] - d(A, B) \\ &= \frac{1}{2}\left[d(x_{n-1}, x_{n+1})\right] \\ &\leq \frac{1}{2}\left[d(x_n, x_{n-1}) + d(x_n, x_{n+1})\right], \end{aligned}$$

and, therefore,

$$\begin{aligned}\psi\left(d(x_{n+1},x_n)\right) &= \psi\left(d(Tx_n,Tx_{n-1})\right)\\ &\leq \psi\left(M(x_n,x_{n-1})-d(A,B)\right)-\phi\left(d(x_n,x_{n-1})\right)\\ &\leq \psi\left(\frac{1}{2}\left[d(x_n,x_{n-1})+d(x_n,x_{n+1})\right]\right)-\phi\left(d(x_n,x_{n-1})\right)\\ &< \psi\left(\frac{1}{2}\left[d(x_n,x_{n-1})+d(x_n,x_{n+1})\right]\right).\end{aligned} \tag{5}$$

Since ψ is nondecreasing

$$d(x_{n+1},x_n) \leq \frac{1}{2}\left[d(x_n,x_{n-1})+d(x_n,x_{n+1})\right],$$

and, consequently,

$$d(x_{n+1},x_n) \leq d(x_n,x_{n-1}). \tag{6}$$

From (5) and (6) we infer that

$$\psi\left(d(x_{n+1},x_n)\right) \leq \psi\left(d(x_n,x_{n-1})\right)-\phi\left(d(x_n,x_{n-1})\right). \tag{7}$$

Summarizing, in any case we get $d(x_{n+1},x_n) \leq d(x_n,x_{n-1})$.

This means that $(d(x_{n+1},x_n))$ is a decreasing sequence of nonnegative real numbers and, consequently, there exists an $r \geq 0$ such that

$$\lim_{n\to\infty} d(x_{n+1},x_n) = r.$$

We shall now prove that $r = 0$.

Notice that the previously discussed Cases 1, 3 and 4 are the only ones possible.

In these cases we obtained (see (3), (4) and (7))

$$\begin{aligned}\psi\left(d(x_{n+1},x_n)\right) &\leq \psi\left(d(x_n,x_{n-1})\right)-\phi\left(d(x_n,x_{n-1})\right)\\ &\leq \psi\left(d(x_n,x_{n-1})\right) \quad \text{for any } n\in\mathbb{N}.\end{aligned}$$

Suppose that $r > 0$.

Taking $n \to \infty$ in the last inequality, and, by using the continuity of ψ, we infer that

$$\psi(r) \leq \psi(r) - \lim_{n\to\infty}\phi\left(d(x_n,x_{n-1})\right) \leq \psi(r),$$

which implies that

$$\lim_{n\to\infty}\phi\left(d(x_n,x_{n-1})\right) = 0. \tag{8}$$

On the other hand, since $r > 0$, we have

$$0 < r \leq d(x_n,x_{n-1}) \quad \text{for any } n\in\mathbb{N},$$

and, since ϕ is nondecreasing and $\phi(t) = 0$ if and only if $t = 0$, we get

$$0 < \phi(r) \leq \phi\left(d(x_n, x_{n-1})\right)) \text{ for any } n \in \mathbb{N}.$$

Therefore, $0 < \phi(r) \leq \lim_{n\to\infty} \phi\left(d(x_n, x_{n-1})\right)$ and this contradicts (8).

Consequently $r = 0$; i.e.,

$$\lim_{n\to\infty} d(x_{n+1}, x_n) = 0. \tag{9}$$

We now prove that (x_n) is a Cauchy sequence

In the contrary case, by using Lemma 2.1 of [6], we can find an $\epsilon > 0$ and subsequences $(x_{n(k)})$ and $(x_{m(k)})$ de (x_n) satisfying:

(i) $n(k) > m(k) \geq k$ for $k \geq 0$.

(ii) $\epsilon \leq d(x_{n(k)}, x_{m(k)})$ and $d(x_{n(k)-1}, x_{m(k)}) < \epsilon$ for $k \geq 0$.

(iii) $$\lim_{k\to\infty} d(x_{n(k)}, x_{m(k)}) = \lim_{k\to\infty} d(x_{n(k)}, x_{m(k)+1}) = \lim_{k\to\infty} d(x_{m(k)}, x_{n(k)+1}) = \lim_{k\to\infty} d(x_{n(k)+1}, x_{m(k)+1}) = \epsilon.$$

Since, for any $k \geq 0$, $d(x_{n(k)+1}, Tx_{n(k)}) = d(A, B)$ and $d(x_{m(k)+1}, Tx_{m(k)} = d(A, B))$, and, as the pair (A, B) has the P-property, we infer that

$$d(x_{n(k)+1}, x_{m(k)+1}) = d(Tx_{n(k)}, Tx_{m(k)}) \text{ for any } k \geq 0.$$

Using the contractive condition, for any $k \geq 0$, we have

$$\begin{aligned} \psi\left(d(x_{n(k)+1}, x_{m(k)+1})\right) &= \psi\left(d(Tx_{n(k)}, Tx_{m(k)})\right) \\ &\leq \psi\left(M(x_{n(k)}, x_{m(k)}) - d(A, B)\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right). \end{aligned} \tag{10}$$

We distinguish four cases.

Case 1: Suppose that $M(x_{n(k)}, x_{m(k)}) = d(x_{n(k)}, x_{m(k)})$. In this case, from the last inequality and by using the fact that ψ is nondecreasing and ϕ is nonnegative, we obtain

$$\begin{aligned} \psi\left(d(x_{n(k)+1}, x_{m(k)+1})\right) &\leq \psi\left(d(x_{n(k)}, x_{m(k)}) - d(A, B)\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right) \\ &\leq \psi\left(d(x_{n(k)}, x_{m(k)})\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right) \\ &\leq \psi\left(d(x_{n(k)}, x_{m(k)})\right). \end{aligned} \tag{11}$$

Case 2: Suppose that $M(x_{n(k)}, x_{m(k)}) = d(x_{n(k)}, Tx_{n(k)})$. In this case,

$$\begin{aligned} M(x_{n(k)}, x_{m(k)}) - d(A, B) &= d(x_{n(k)}, Tx_{n(k)}) - d(A, B) \\ &\leq d(x_{n(k)}, x_{n(k)+1}) + d(x_{n(k)+1}, Tx_{n(k)}) - d(A, B) \\ &= d(x_{n(k)}, x_{n(k)+1}), \end{aligned}$$

and inequality (10) takes the form

$$\begin{aligned} \psi\left(d(x_{n(k)+1}, x_{m(k)+1})\right) &\leq \psi\left(d(x_{n(k)}, x_{n(k)+1})\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right) \\ &\leq \psi\left(d(x_{n(k)}, x_{n(k)+1})\right). \end{aligned} \tag{12}$$

Case 3: Suppose that $M(x_{n(k)}, x_{m(k)}) = d(x_{m(k)}, Tx_{m(k)})$. Using an argument similar to that in Case 2, we have

$$M(x_{n(k)}, x_{m(k)}) - d(A, B) \leq d(x_{m(k)}, x_{m(k)+1}),$$

and, from (10), we have

$$\begin{aligned} \psi\left(d(x_{n(k)+1}, x_{m(k)+1})\right) &\leq \psi\left(d(x_{m(k)}, x_{m(k)+1})\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right) \\ &\leq \psi\left(d(x_{m(k)}, x_{m(k)+1})\right). \end{aligned} \tag{13}$$

Case 4: Suppose that $M(x_{n(k)}, x_{m(k)}) = \frac{1}{2}\left[d(x_{n(k)}, Tx_{m(k)}) + d(x_{m(k)}, Tx_{n(k)})\right]$. In this case, we have that

$$\begin{aligned} & M(x_{n(k)}, x_{m(k)}) - d(A, B) \\ &= \frac{1}{2}\left[d(x_{n(k)}, Tx_{m(k)}) + d(x_{m(k)}, Tx_{n(k)})\right] - d(A, B) \\ &\leq \frac{1}{2}\left[d(x_{n(k)}, x_{m(k)+1}) + d(x_{m(k)+1}, Tx_{m(k)})\right. \\ &\qquad \left. + d(x_{m(k)}, x_{n(k)+1}) + d(x_{n(k)+1}), Tx_{n(k)})\right] - d(A, B) \\ &= \frac{1}{2}\left[d(x_{n(k)}, x_{m(k)+1}) + d(A, B) + d(x_{m(k)}, x_{n(k)+1}) + d(A, B)\right] - d(A, B) \\ &= \frac{1}{2}\left[d(x_{n(k)}, x_{m(k)+1}) + d(x_{m(k)}, x_{n(k)+1})\right], \end{aligned}$$

and, consequently, from (10) we infer that

$$\begin{aligned} & \psi\left(d(x_{n(k)+1}, x_{m(k)+1})\right) \\ &\leq \psi\left(\frac{1}{2}\left[d(x_{n(k)}, x_{m(k)+1}) + d(x_{m(k)}, x_{n(k)+1})\right]\right) - \phi\left(d(x_{n(k)}, x_{m(k)})\right). \end{aligned} \tag{14}$$

Now, put

$$\begin{aligned} C &= \{k \geq 0 \colon k \text{ satisfies Case 2}\}, \\ D &= \{k \geq 0 \colon k \text{ satisfies Case 3}\}, \quad \text{and} \\ E &= \{k \geq 0 \colon k \text{ satisfies Case 4}\}. \end{aligned}$$

If $\operatorname{Card} C = \infty$ or $\operatorname{Card} D = \infty$ then, taking $k \to \infty$ in (12) or (13) and, taking into account the properties of the subsequences $(x_{n(k)})$ and $(x_{m(k)})$, (9) and the continuity of ψ, we get

$$0 < \psi(\epsilon) = \psi\left(\lim_{k\to\infty} d(x_{n(k)}, x_{n(k)+1})\right) = \psi(0) = 0$$

or

$$0 < \psi(\epsilon) = \psi\left(\lim_{k\to\infty} d(x_{m(k)}, x_{m(k)+1})\right) = \psi(0) = 0$$

which is a contradiction.

Therefore $\operatorname{Card} C < \infty$ and $\operatorname{Card} D < \infty$.

If $\operatorname{Card} E = \infty$ then taking $k \to \infty$ in (14), we have

$$0 < \psi(\epsilon) \leq \psi\left(\frac{1}{2}(\epsilon+\epsilon)\right) - \lim_{k\to\infty} \phi\big(d(x_{n(k)}, x_{m(k)}\big) \leq \psi(\epsilon),$$

which implies that

$$\lim_{k\to\infty} \phi\big(d(x_{n(k)}, x_{m(k)})\big) = 0. \tag{15}$$

Since $0 < \epsilon \leq d(x_{n(k)}, x_{m(k)})$ for any $k \geq 0$ and ϕ is nondecreasing, we get

$$0 < \phi(\epsilon) \leq \phi\big(d(x_{n(k)}, x_{m(k)})\big), \quad \text{for any } k \geq 0.$$

Therefore $\operatorname{Card} E < \infty$.
which contradicts (15).
This means that, for all $k \geq 0$ except for a finite number, Case 1 is satisfied.

Taking $k \to \infty$ in (11) we obtain

$$0 < \psi(\epsilon) \leq \psi(\epsilon) - \lim_{k\to\infty} \phi\left(d(x_{n(k)}, x_{m(k)})\right) \leq \psi(\epsilon),$$

which implies that

$$\lim_{k\to\infty} \phi\left(d(x_{n(k)}, x_{m(k)})\right) = 0. \tag{16}$$

But, since

$$0 < \epsilon \leq d(x_{n(k)}, x_{m(k)}) \quad \text{for all } k \geq 0,$$

by using the nondecreasing character of ϕ, we have

$$0 < \phi(\epsilon) \leq \phi\left(d(x_{n(k)}, x_{m(k)})\right) \quad \text{for all } k \geq 0,$$

and, from this fact, we infer that

$$0 < \phi(\epsilon) \leq \lim_{k\to\infty} \phi\left(d(x_{n(k)}, x_{m(k)})\right),$$

which contradicts (16).

Therefore, $\{x_n\}$ is a Cauchy sequence.

Since $\{x_n\} \subset A$ and A is a closed subset of the complete metric space (X, d), we can find an $x^* \in A$ such that $x_n \to x^*$.

Now, since T is continuous, $Tx_n \to Tx^*$.

Therefore, we have that $d(x_{n+1}, Tx_n) \to d(x^*, Tx^*)$. Taking into account that the sequence $d(x_{n+1}, Tx_n)$ is a constant sequence with value $d(A, B)$, we deduce that

$$d(x^*, Tx^*) = d(A, B).$$

This means that x^* is a best proximity point of T.

This proves the existence part of our theorem.

To prove uniqueness, suppose that x_1 and x_2 are two best proximity points of T and $x_1 \neq x_2$. This means that

$$d(x_i, Tx_i) = d(A, B) \text{ for } i = 1, 2.$$

By using the P-property of the pair (A, B), we have

$$d(x_1, x_2) = d(Tx_1, Tx_2).$$

Now, taking into account the contractive condition, we get

$$\psi\left(d(x_1, x_2)\right) = \psi\left(d(Tx_1, Tx_2)\right) \leq \psi\left(M(x_1, x_2) - d(A, B)\right) - \phi\left(d(x_1, x_2)\right). \quad (17)$$

There are four cases.

Case 1: Suppose that $M(x_1, x_2) = d(x_1, x_2)$. In this case inequality (17) takes the form

$$\psi\left(d(x_1, x_2)\right) \leq \psi\left(d(x_1, x_2) - d(A, B)\right) - \phi\left(d(x_1, x_2)\right),$$

and, since $x_1 \neq x_2$ and $\phi(t) = 0$ if and only if $t = 0$, we have that $\phi\left(d(x_1, x_2)\right) > 0$. Consequently, the last inequality gives us

$$\begin{aligned} \psi\left(d(x_1, x_2)\right) &\leq \psi\left(d(x_1, x_2) - d(A, B)\right) - \phi\left(d(x_1, x_2)\right) \\ &< \psi\left(d(x_1, x_2) - d(A, B)\right) \\ &\leq \psi\left(d(x_1, x_2)\right), \end{aligned}$$

and this is a contradiction.

Case 2: Suppose that $M(x_1, x_2) = d(x_1, Tx_1)$. Since $d(x_1, Tx_1) = d(A, B)$, the inequality (17) takes the form

$$\begin{aligned} \psi\left(d(x_1, x_2)\right) &\leq \psi\left(d(x_1, Tx_1) - d(A, B)\right) - \phi\left(d(x_1, x_2)\right) \\ &= \psi\left(d(A, B) - d(A, B)\right) - \phi\left(d(x_1, x_2)\right) \\ &= \psi(0) - \phi\left(d(x_1, x_2)\right). \end{aligned}$$

As $\psi(0) = 0$ and $x_1 \neq x_2$ and, consequently, $\phi\left(d(x_1, x_2)\right) > 0$, we have

$$\psi\left(d(x_1, x_2)\right) \leq -\phi\left(d(x_1, x_2)\right) < 0,$$

which is a contradiction.

Case 3: Suppose that $M(x_1, x_2) = d(x_2, Tx_2)$. Using the same argument as in Case 2, we obtain

$$\psi\left(d(x_1, x_2)\right) \leq -\phi\left(d(x_1, x_2)\right) < 0,$$

which is a contradiction.

Case 4: Suppose that $M(x_1, x_2) = \frac{1}{2}[d(x_1, Tx_2) + d(x_2, Tx_1)]$. Since

$$\begin{aligned} & M(x_1, x_2) - d(A, B) \\ &= \frac{1}{2}[d(x_1, Tx_2) + d(x_2, Tx_1)] - d(A, B) \\ &\leq \frac{1}{2}[d(x_1, Tx_1) + d(Tx_1, Tx_2) + d(x_2, Tx_2 + d(Tx_1, Tx_2))] - d(A, B) \\ &= \frac{1}{2}[d(A, B) + 2d(Tx_1, Tx_2) + d(A, B)] - d(A, B) \\ &= d(Tx_1, Tx_2), \end{aligned}$$

from (17), we get

$$\begin{aligned} & \psi(d(x_1, x_2)) \\ &= \psi(d(Tx_1, Tx_2)) \leq \psi(d(Tx_1, Tx_2)) - \phi(d(x_1, x_2)) \leq \psi(d(Tx_1, Tx_2)). \end{aligned}$$

Consequently, $\phi(d(x_1, x_2)) = 0$.
This is a contradiction since $x_1 \neq x_2$.

Therefore $x_1 = x_2$. □

In what follows we will present some particular cases and their consequences.

When the mapping ψ in Theorem 3.1 is the identity, we obtain the following corollary.

Corollary 3.2. *Let (A, B) a pair of nonempty closed subsets of a complete metric space (X, d) such that $A_0 \neq \emptyset$. Let $T\colon A \to B$ be a continuous mapping satisfying that $T(A_0) \subset B_0$ and such that*

$$d(Tx, Ty) \leq M(x, y) - d(A, B) - \phi(d(x, y)) \quad \text{for any } x, y \in A.$$

Suppose that the pair (A, B) has the P-property. Then there exists a unique $x^* \in A$ such that $d(x^*, Tx^*) = d(A, B)$.

When $d(A, B) = 0$, Corollary 3.2 takes the following form.

Corollary 3.3. *Let (A, B) be a pair of nonempty closed subsets of a complete metric space (X, d) such that $A_0 \neq \emptyset$ and $d(A, B) = 0$. Let $T\colon A \to B$ be a continuous mapping such that $T(A_0) \subset B_0$ and satisfying*

$$d(Tx, Ty) \leq M(x, y) - \phi(d(x, y)) \quad \text{for any } x, y \in A.$$

Then there exists a unique $x^* \in A$ such that $Tx^* = x^*$.

Proof. Notice that, if $d(A, B) = 0$, then (A, B) has the P-property. □

The condition

$$d(Tx, Ty) \leq d(x, y) - \phi(d(x, y)), \text{ for any } x, y \in A$$

implies that

$$d(Tx,Ty) \leq d(x,y) - \phi\left(d(x,y)\right) \leq d(x,y) \text{ for any } x,y \in A.$$

Thus T is continuous and satisfies

$$d(Tx,Ty) \leq d(x,y) - \phi\left(d(x,y)\right) \leq M(x,y) - \phi\left(d(x,y)\right) \text{ for any } x,y \in A.$$

By Theorem 3.1 we obtain the desired result.
As a consequence of Corollary 3.3 we have the following result.

Corollary 3.4. *Let (A,B) be a pair of nonempty closed subsets of a complete metric space (X,d) such that $A_0 \neq$ and $d(A,B) = 0$. Let $T\colon A \to B$ be a mapping such that $T(A_0) \subset B_0$ and satisfies*

$$d(Tx,Ty) \leq d(x,y) - \phi\left(d(x,y)\right), \text{ for any } x,y \in A.$$

Then there exists a unique $x^* \in A$ such that $Tx^* = x^*$.

Corollary 3.4 is Theorem 3.1 of [12] for the particular case $d(A,B) = 0$.

The following result presents a best proximity point theorem for a mapping satisfying a Kannan type contractive condition.

Corollary 3.5. *Let (A,B) be a pair of nonempty closed subsets of a complete metric space (X,d) such that $A_0 \neq \emptyset$. Let $T\colon A \to B$ be a continuous mapping satisfying $T(A_0) \subset B_0$ and such that*

$$\begin{aligned} &\psi\left(d(Tx,Ty)\right) \\ &\leq \psi\left(\frac{1}{2}[d(x,Tx) + d(y,Ty)] - d(A,B)\right) - \phi\left(d(x,y)\right), \text{ for any } x,y \in A, \end{aligned}$$

where $\psi\colon [0,\infty) \to [0,\infty)$ is a continuous and nondecreasing function such that $\psi(t) = 0$ if and only if $t = 0$ and $\phi\colon [0,\infty) \to [0,\infty)$ is a nondecreasing function such that $\phi(t) = 0$ if and only if $t = 0$.

Suppose that the pair (A,B) has the P-property. Then there exists a unique $x^* \in A$ such that $d(x^*,Tx^*) = d(A,B)$.

Proof. First, notice that

$$\frac{1}{2}[d(x,Tx) + d(y,Ty)] \geq \frac{1}{2}[d(A,B) + d(A,B)] = d(A,B),$$

and, consequently, the contractive condition is well defined.

On the other hand, since

$$\frac{1}{2}[d(x,Tx) + d(y,Ty)] \leq \frac{1}{2}[M(x,y) + M(x,y)] = M(x,y)$$

we have, by virtue of the nondecreasing character of ψ that, for any $x, y \in A$

$$\begin{aligned}\psi\left(d(Tx,Ty)\right) &\leq \psi\left(\frac{1}{2}[d(x,Tx)+d(y,Ty)]-d(A,B)\right)-\phi\left(d(x,y)\right)\\ &\leq \psi\left(M(x,y)-d(A,B)\right)-\phi\left(d(x,y)\right),\end{aligned}$$

and the desired result is obtained by applying Theorem 3.1. □

The following consequence is a best proximity point theorem for mappings satisfying a condition of integral type in the case $d(A,B)=0$.

Corollary 3.6. *Let (A,B) be a pair of nonempty closed subsets of a complete metric space (X,d) such that $d(A,B)=0$ and $A_0 \neq \emptyset$. Let $T\colon A \to B$ be a continuous mapping such that $T(A_0) \subset B_0$ and suppose that there exists a $k \in [0,1)$ such that*

$$\int_0^{d(Tx,Ty)} \varphi(t)\,dt \leq k \int_0^{M(x,y)} \varphi(t)\,dt \quad \text{for any } x, y \in A,$$

where $\varphi\colon \mathbb{R}_+ \to \mathbb{R}_+$ is a Lebesgue-integrable mapping such that $\int_0^\epsilon \varphi(t)\,dt > 0$ for $\epsilon > 0$.

Then there exists a unique $x^* \in A$ such that $Tx^* = x^*$.

Proof. First notice that, if $d(A,B)=0$, then it is easily seen that (A,B) has the P-property. On the other hand, it is easily seen that the function $\psi\colon [0,\infty) \to [0,\infty)$ defined by

$$\psi(t) = \int_0^t \varphi(s)\,ds$$

is continuous and nondecreasing and, since $\int_0^\epsilon \varphi(t)\,dt > 0$ for each $\epsilon > 0$, $\psi(t) = 0$ if and only if $t = 0$.

Let ϕ be the function defined by $\phi(t) = (1-k)\psi(t)$ on $[0,\infty)$. Therefore our contractive mapping can be rewritten of the following way

$$\begin{aligned}\psi\left(d(Tx,Ty)\right) &\leq k\psi\left(M(x,y)\right)\\ &= \psi\left(M(x,y)\right)-(1-k)\psi\left(M(x,y)\right)\\ &= \psi\left(M(x,y)\right)-\phi\left(M(x,y)\right), \quad \text{for any } x, y \in A.\end{aligned}$$

By using the nondecreasing character of ϕ, the last inequality implies that

$$\psi\left(d(Tx,Ty)\right) \leq \psi\left(M(x,y)\right)-\phi\left(d(x,y)\right), \text{ for any } x, y \in A.$$

By applying Theorem 3.1, we obtain the desired result. □

When $A = B$ Theorem 3.1 has, as consequences, the following fixed point theorems.

Corollary 3.7. *Let (X,d) be a complete metric space and let $T\colon X \to X$ be a mapping satisfying*

$$\psi\left(d(Tx,Ty)\right) \leq \psi\left(M(x,y)\right) - \phi\left(d(x,y)\right), \quad \text{for any } x,y \in X,$$

where $\psi\colon [0,\infty) \to [0,\infty)$ is a continuous and nondecreasing function such that $\psi(t)=0$ if and only if $t=0$ and $\phi\colon [0,\infty) \to [0,\infty)$ is a nondecreasing function such that $\phi(t)=0$ if and only if $t=0$. Then T has a unique fixed point.

Corollary 3.7 has as a consequence, the following result.

Corollary 3.8. *Let (X,d) be a complete metric space and let $T\colon X \to X$ be a mapping satisfying*

$$\psi\left(d(Tx,Ty)\right) \leq \psi\left(d(x,y)\right) - \phi\left(d(x,y)\right), \quad \text{for any } x,y \in X,$$

where ψ and ϕ satisfy the same conditions as those in Corollary 3.7. Then T has a unique fixed point.

Proof. Notice that, since ψ is nondecreasing, for any $x,y \in X$

$$\psi\left(d(Tx,Ty)\right) \leq \psi\left(d(x,y)\right) - \phi\left(d(x,y)\right) \leq \psi\left(M(x,y)\right) - \phi\left(d(x,y)\right).$$

Moreover, since $\psi\left(d(Tx,Ty)\right) \leq \psi\left(d(x,y)\right)$ for any $x,y \in X$ and ψ is nondecreasing, $d(Tx,Ty) \leq d(x,y)$ for any $x,y \in X$ and thus T is continuous. Finally, Corollary 3.7 give us the desired result. □

Corollary 3.8 improves Theorem 2.1 of [4] since we do not assume the continuity of ϕ.

If we put $\psi = 1_X$(identity mapping) in Corollary 3.8 then we obtain the following result.

Corollary 3.9. *Let (X,d) be a complete metric space and $T\colon X \to X$ a mapping satisfying*

$$d(Tx,Ty) \leq d(x,y) - \phi\left(d(x,y)\right), \quad \text{for any } x,y \in X$$

where ϕ satisfies the same conditions as in Corollary 3.7. Then T has a unique fixed point.

Corollary 3.9 improves Theorem 1 of [10] because we do not impose the continuity of ϕ.

A consequence of Corollary 3.6 is the following fixed point theorem with a condition of integral type.

Corollary 3.10. *Let (X,d) be a complete metric space and $k \in [0,1)$ and let $T\colon X \to X$ be a mapping such that for each $x,y \in X$*

$$\int_0^{d(Tx,Ty)} \varphi(t)\,dt \leq k \int_0^{M(x,y)} \varphi(t)\,dt,$$

where $\varphi\colon \mathbb{R}_+ \to \mathbb{R}_+$ *is a Lebesgue-integrable mapping such that* $\int_0^\epsilon \varphi(t)\,dt > 0$ *for* $\epsilon > 0$. *Then* T *has a unique fixed point.*

Since the contractive condition

$$\int_0^{d(Tx,Ty)} \varphi(t)\,dt \leq k \int_0^{d(x,y)} \varphi(t)\,dt \quad \text{for any } x, y \in X$$

implies the one appearing in Corollary 3.10, we have the following result.

Corollary 3.11. *Let* (X, d) *be a complete metric space and* $k \in [0, 1)$ *and* $T\colon X \to X$ *a continuous mapping satisfying*

$$\int_0^{d(Tx,Ty)} \varphi(t)\,dt \leq k \int_0^{d(x,y)} \varphi(t)\,dt \quad \textit{for any } x, y \in X$$

where φ *satisfies the same conditions that in Corollary* 3.10. *Then* T *has a unique fixed point.*

Corollary 3.11 is the main result of [3] when T is continuous.

The following example shows that the P-property condition cannot be relaxed from Theorem 3.1.

Example 3.12. Consider $X = \mathbb{R}$ with the usual metric and $A = \{8, -8\}$ and $B = \{1, -1\}$.

Obviously, A and B are closed subsets of X with $A_0 = A$, $B_0 = B$ and $d(A, B) = 7$. Let $T\colon A \to B$ be the mapping defined by $T(8) = -1$ and $T(-8) = 1$. It is easily seen that

$$d(Tx, Ty) \leq \frac{1}{2}\,(d(x, y) - d(A, B)) \quad \text{for } x, y \in A,$$

and, thus, T satisfies the contractive condition appearing in Theorem 3.1. On the other hand, since $d(x, Tx) = 9 > d(A, B) = 7$ for any $x, y \in A$, T does not have a best proximity point.

Notice that the pair (A, B) does not have the P-property

4. Examples.

In order to illustrate our results, we present some examples.

Example 4.1. Consider $X = \mathbb{R}^2$ with the usual metric and the closed subsets

$$A = [0, \infty) \times \{0\} \quad \text{and} \quad B = (-\infty, 0] \times \{0\}.$$

Obviously, $d(A, B) = 0$ and, consequently, the pair (A, B) has the P-property. Notice that $A_0 = \{(0, 0)\}$ and $B_0 = \{(0, 0)\}$.

Let $T\colon A \to B$ be the mapping defined by

$$T(x, 0) = (-\log(1 + x), 0).$$

Obviously $T(A_0) \subset B_0$.

In the sequel, we will prove that T satisfies the contractive condition of Theorem 3.1.

In fact, for $(x,0),(y,0) \in A$, we have

$$\begin{aligned} d(T(x,0),T(y,0)) &= d\left((-\log(1+x),0),(-\log(1+y),0)\right) \\ &= |\log(1+x)-\log(1+y)| \\ &= \left|\log\left(\frac{1+x}{1+y}\right)\right|. \end{aligned}$$

We claim that $\left|\log\left(\frac{1+x}{1+y}\right)\right| \leq \log(1+|x-y|)$.

Suppose that $x > y$(the same argument works for $y > x$).

Since $\phi(t) = \log(1+t)$ is strictly increasing in $[0,\infty)$, we have

$$\begin{aligned} \log\left(\frac{1+x}{1+y}\right) &= \log\left(\frac{1+y+x-y}{1+y}\right) = \log\left(1+\frac{x-y}{1+y}\right) \\ &\leq \log(1+x-y) = \log(1+|x-y|). \end{aligned}$$

This prove our claim.

Therefore, for $(x,0),(y,0) \in A$, we have

$$\begin{aligned} & d\left(T(x,0),T(y,0)\right) \\ =\ & \left|\log\left(\frac{1+x}{1+y}\right)\right| \\ \leq\ & \log(1+|x-y|) \\ =\ & \log\left(1+d\left((x,0),(y,0)\right)\right) \\ =\ & d\left((x,0),(y,0)\right) - \left(d\left((x,0),(y,0)\right) - \log\left(1+d\left((x,0),(y,0)\right)\right)\right). \end{aligned}$$

In our case, $\psi = 1_{[0,\infty)}$(identity mapping) and ϕ is given by $\phi(t) = t - \log(1+t)$ which is nondecreasing and satisfies that $\phi(t) = 0$ if and only if $t = 0$. T is obviously continuous and, therefore, the assumptions of Theorem 3.1 are satisfied. Consequently, by Theorem 3.1 there exists a unique $(x^*,0) \in A$ such that

$$\left(d\left((x^*,0),T(x^*,0)\right) = 0 = d(A,B).\right.$$

Notice that $(x^*,0) \in A$ is the point $(0,0)$.

The following example shows that the condition A and B are closed subsets of (X,d) is not a necessary condition for the existence of a unique best proximity point.

Example 4.2. Consider $X = \mathbb{R}^2$ with the usual metric and the subsets

$$A = [0,\infty) \times \{0\} \quad \text{and} \quad B = \left(-\frac{\pi}{2},0\right] \times \{0\}.$$

Notice that B is not a closed subset of X.

It is easily seen that $d(A,B)=0$, $A_0=\{(0,0)\}$ and $B_0=\{(0,0)\}$.

Since $d(A,B)=0$ then the pair (A,B) has the P-property.

Let $T\colon A\to B$ the mapping defined by

$$T(x,0)=(-\arctan x,0).$$

Obviously, $T(A_0)\subset B_0$.

Now, we check that T satisfies the contractive condition of Theorem 3.1.

In fact, for $(x,0),(y,0)\in A$, we have

$$d\left(T(x,0),T(y,0)\right)=d\left((-\arctan x,0),(-\arctan y,0)\right)=\left|\arctan x-\arctan y\right|.$$

We claim that $|\arctan x-\arctan y|\leq \arctan|x-y|$.

In fact, suppose that $x>y$(the same reasoning works for $y>x$).

Put $\arctan x=\alpha$ and $\arctan y=\beta$ (notice that $\alpha>\beta$ since the function $\arctan t$ for $t\geq 0$ is strictly increasing).

Now, since

$$\tan(\alpha-\beta)=\frac{\tan\alpha-\tan\beta}{1+\tan\alpha\tan\beta},$$

and, since $\alpha,\beta\in\left[0,\frac{\pi}{2}\right)$, we have that $\tan\alpha,\tan\beta\in[0,\infty)$. Thus, from the last expression it follows that

$$\tan(\alpha-\beta)\leq\tan\alpha-\tan\beta.$$

Applying ϕ to the last inequality (where $\phi(t)=\arctan t$) and, by using the nondecreasing character of ϕ, we infer that

$$\alpha-\beta\leq\arctan(\tan\alpha-\tan\beta).$$

This means that

$$\arctan x-\arctan y=\alpha-\beta\leq\arctan(x-y).$$

This proves our claim.

Therefore, we get, for $(x,0),(y,0)\in A$

$$\begin{aligned}
& d\left(T(x,0),T(y,0)\right)\\
=\ & \left|\arctan x-\arctan y\right|\\
\leq\ & \arctan|x-y|\\
=\ & \arctan\left(d\left((x,0),(y,0)\right)\right)\\
=\ & d\left((x,0),(y,0)\right)-\left(d\left((x,0),(y,0)\right)-\arctan\left(d\left((x,0),(y,0)\right)\right)\right).
\end{aligned}$$

In our case, $\psi = 1_{[0,\infty)}$(identity mapping) and ϕ is given by $\phi(t) = t - \arctan t$. It is easily seen that ϕ is nondecreasing and $\phi(t) = 0$ if and only if $t = 0$. Obviously, T is continuous, and satisfies assumptions of Theorem 3.1.

Notice that $(0, 0) \in A$ is the unique best proximity point of T, since

$$d\left((0,0), T(0,0)\right) = 0 = d(A, B).$$

Therefore, the conclusion of Theorem 3.1 is satisfied, but B is not a closed subset.

References

[1] M. A. Al-Thagafi, N. Shahzad: Convergence and existence results for best proximity points, Nonlinear Anal. 70 (2009) 3665–3671.

[2] Ya. I. Alber, S. Guerre-Delabriere: Principle of weakly contractive maps in Hilbert spaces, in: New Results in Operator Theory and its Applications, I. Gohberg et al. (ed.), Operator Theory: Advances and Applications 98, Birkhäuser, Basel (1997) 7–22.

[3] A. Branciari: A fixed point theorem for mappings satisfying a general contractive condition of integral type, Int. J. Math. Math. Sci. 29 (2002) 531–536.

[4] P. N. Dutta, B. S. Choudhury: A generalisation of contraction principle in metric spaces, Fixed Point Theory Appl. (2008), Article ID 406368, 8 pp.

[5] A. A. Eldred, P. Veeramani: Existence and convergence of best proximity points, J. Math. Anal. Appl. 323 (2006) 1001–1006.

[6] M. Jleli, V. C. Rajić, B. Samet, C. Vetro: Fixed point theorems on ordered metric spaces and applications to nonlinear elastic beam equations, J. Fixed Point Theory Appl. 12 (2012) 175–192.

[7] W. A. Kirk, S. Reich, P. Veeramani: Proximinal retracts and best proximity pair theorems, Numer. Funct. Anal. Optim. 24 (2003) 851–862.

[8] W. A. Kirk, B. Sims (Eds.): Handbook of Metric Fixed Point Theory, Kluwer, Dordrecht (2001).

[9] B. E. Rhoades: A comparison of various definitions of contraction mappings, Trans. Amer. Math. Soc. 226 (1997) 257–290.

[10] B. E. Rhoades: Some theorems on weakly contractive maps, Nonlinear Anal. (2001) 2683–2693.

[11] S. Sadiq Basha, P. Veeramani: Best proximity pair theorems for multifunctions with open fibres, J. Approx. Theory 103 (2000) 119–129.

[12] V. Sankar Raj: A best proximity theorem for weakly contractive non-self mappings, Nonlinear Anal. 74 (2011) 4804–4808.

[13] V. Sankar Raj: Banach's contraction principle for non-selt mappings, preprint.