

# An Existence Result for Quasi-Equilibrium Problems

**D. Aussel**

*Laboratoire PROMES, Université de Perpignan, Tecnosud,  
Rambla de la Thermodynamique, 66100 Perpignan, France  
aussel@univ-perp.fr*

**J. Cotrina**

*Centro de Investigación de la Universidad del Pacífico,  
Jr. Sanchez Cerro 2141 Jesús María, Lima 11, Perú  
cotrina\_je@up.edu.pe*

**A. N. Iusem**

*Instituto de Matemática Pura e Aplicada (IMPA),  
Estrada Dona Castorina 110, Rio de Janeiro, RJ, CEP 22460-320, Brazil  
iusp@impa.br*

Received: January 3, 2014

Revised manuscript received: April 7, 2015

Accepted: April 10, 2015

Recently, it has been showed in Castellani-Giuli [13], that the proof of the existence result for quasimonotone Stampacchia variational inequalities developed in Aussel-Hadjisavvas [6] can be adapted to the case of equilibrium problems. This proof was based on KKM techniques. In this paper we define and study the so-called quasi-equilibrium problem, that is an equilibrium problem with a constraint set depending on the current point. Our main contribution consists of an existence result combining fixed point techniques with stability analysis of perturbed equilibrium problems.

## 1. Introduction

The equilibrium problem, (EP) in short, is defined as follows. Given a real Banach space  $X$ , a nonempty subset  $K$  of  $X$  and a bifunction  $f : X \times X \rightarrow \mathbb{R}$ , (EP) consists of

$$(EP) \quad \text{find } x \in K \text{ such that } f(x, y) \geq 0 \quad \forall y \in K.$$

Problem (EP) has been extensively studied in recent years (see e.g. [19, 20, 24, 25, 26]). A recurrent theme in the analysis of the conditions for the existence of solutions of (EP) is the connection between them and the solutions of the so-called *Convex Feasibility Problem*, to be denoted (CFP), which turns out to be convex under suitable conditions on  $f$  and which corresponds to a sort of dual formulation of (EP),

$$(CFP) \quad \text{find } x \in K \text{ such that } f(y, x) \leq 0 \quad \forall y \in K.$$

It was proved in [26] that if  $f$  is lower semicontinuous in the first argument, convex in the second one and it vanishes on the diagonal of  $K \times K$ , then every solution of (CFP) is a solution of (EP), and moreover both solution sets trivially coincide under pseudomonotonicity of  $f$ . Bianchi *et al.* in [10] extended this inclusion under a weak continuity property of the bifunction, and they obtained an existence result for (EP), adapting the existence result for variational inequality proposed by Aussel *et al.* in [6].

The classical example of equilibrium problem is the variational inequality problem, which is defined as follows: a Stampacchia variational inequality problem (VIP) is formulated as

$$(VIP) \quad \begin{array}{l} \text{find } x \in K \text{ such that there exists } x^* \in T(x) \\ \text{with } \langle x^*, y - x \rangle \geq 0, \forall y \in K, \end{array}$$

where  $T : X \rightarrow 2^{X^*}$  is a set-valued operator,  $X^*$  is the dual space of  $X$  and  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $X$  and  $X^*$ . So, if  $T$  has compact values, and we define the *representative bifunction*  $f_T$  of  $T$  by

$$f_T(x, y) = \sup_{x^* \in T(x)} \langle x^*, y - x \rangle,$$

it follows that every solution of the Equilibrium Problem associated to  $f_T$  and  $K$  is a solution of the Variational Inequality Problem associated to  $T$  and  $K$ , and conversely. Now, the CFP associated to  $f_T$  is equivalent to

$$\text{find } x \in K \text{ such that } \langle y^*, y - x \rangle \geq 0, \forall y \in K, y^* \in T(y)$$

which is known as the Minty variational inequality problem (or dual variational inequality problem).

We denote by  $S(T, K)$  and  $M(T, K)$  the solution sets of the Stampacchia and Minty variational inequality problems respectively.

Recently, it was showed in [11, 15, 23, 27] that any equilibrium problem for which  $f$  is lower semicontinuous in the first argument, convex in the second one, monotone, and vanishes on the diagonal of  $K \times K$ , can be reformulated as a variational inequality. Castellani *et al.* in [12] extended these results to the pseudomonotone and quasimonotone case. In this work we extend them to the quasi-equilibrium problem.

We consider next the problem which is our main object of interest in this paper. The quasi-equilibrium problem, (QEP) in short, is defined as follows. Given a set-valued map  $K : X \rightarrow 2^X$  and a bifunction  $f : X \times X \rightarrow \mathbb{R}$ , (QEP) consists of

$$(QEP) \quad \text{find } x \in K(x) \text{ such that } f(x, y) \geq 0, \forall y \in K(x).$$

The associated Minty quasi-equilibrium problem (to be denoted QMEP), consists of

$$(QMEP) \quad \text{find } x \in K(x) \text{ such that } f(y, x) \leq 0, \forall y \in K(x).$$

We denote by  $\text{QEP}(f, K)$  and  $\text{QMEP}(f, K)$  the sets of solutions of the quasi-equilibrium and Minty quasi-equilibrium problem, respectively.

One of the main reasons for studying quasi-equilibrium problems lies in the relation between them and quasi-variational inequalities (see Section 3.2), which mirrors the well known relation between equilibrium problems and variational inequalities. Quasi-variational inequalities are themselves relevant because they encompass certain problems of interest in various fields of application, which do not fall within the scope of variational inequalities. Perhaps the most important instance of this situation is the generalized Nash equilibrium problem, which models a large number of real life problems in Economics and other areas (see e.g. [17], [30] and references therein). The reduction of generalized Nash equilibrium problems to quasi-variational inequalities has been analyzed, e.g., in [2] and [30]. Existence results for problem  $(QEP)$  can be found in [22] and references therein.

The paper is organized as follows. First in Section 2 we describe the main notation and give some preliminary results and comments concerning the mainly used concepts, in particular the upper sign property of a bifunction. Section 3 is devoted to the study of the relationship between  $(QEP)$  and  $(QMEP)$  on one side and between  $(EP)$  and a variational reformulation on the other hand. Finally in Section 4 we prove an existence result for quasi-equilibrium problem  $(QEP)$  using fixed point techniques.

## 2. Preliminaries

All along the paper,  $X$  stands for a real Banach space,  $X^*$  for its topological dual and  $\langle \cdot, \cdot \rangle$  for the associated duality product. The subsets  $S^*$  and  $B^*$  stand respectively for the unit sphere and unit ball  $S^* = \{x^* \in X^*; \|x^*\| = 1\}$  and  $B^* = \{x^* \in X^*; \|x^*\| \leq 1\}$  of the dual space  $X^*$ .

First, recall that a bifunction  $f : X \times X \rightarrow \mathbb{R}$  is said to be

- quasimonotone on a subset  $K$  if, for all  $x, y \in K$ ,

$$f(x, y) > 0 \Rightarrow f(y, x) \leq 0,$$

- properly quasimonotone on a subset  $K$  if, for all  $x_1, x_2, \dots, x_n \in K$ , and all  $x \in \text{co}(\{x_1, x_2, \dots, x_n\})$ , there exists  $i \in \{1, 2, \dots, n\}$  such that

$$f(x_i, x) \leq 0,$$

- pseudomonotone on a subset  $K$  if, for all  $x, y \in K$ ,

$$f(x, y) \geq 0 \Rightarrow f(y, x) \leq 0.$$

Clearly, pseudomonotonicity of  $f$  implies proper quasimonotonicity of  $f$ . Note that nevertheless no relationship exists between quasimonotonicity and proper quasimonotonicity of bifunctions, see e.g. counter-examples in [9].

Let us recall that a function  $h : X \rightarrow \mathbb{R}$  is said to be:

- *quasiconvex* on a convex subset  $K$  if, for all  $x, y \in K$  and all  $z \in [x, y]$ ,

$$h(z) \leq \max \{h(x), h(y)\}$$

- *semistrictly quasiconvex* on a convex subset  $K$  if  $h$  is quasiconvex on  $K$  and

$$h(x) < h(y) \Rightarrow h(z) < h(y), \quad \forall z \in [x, y[$$

for all  $x, y \in K$ .

In the sequel, we will sometimes use the following assumptions on the considered bifunctions:

- (H<sub>1</sub>)  $f(x, x) = 0$  for all  $x \in K$ ,
- (H<sub>2</sub>)  $f(x, \cdot)$  is semistrictly quasiconvex for all  $x \in K$ ,
- (H<sub>3</sub>)  $f(x, \cdot)$  is lower semicontinuous for all  $x \in K$ .

Our forthcoming stability analysis will be based on the following weak continuity property for bifunctions:

**Definition 2.1.** A bifunction  $f$  is said to have the upper sign property at  $x \in K$  if for every  $y \in K$  the following implication holds:

$$(f(x_t, x) \leq 0, \quad \forall t \in ]0, 1[) \Rightarrow f(x, y) \geq 0, \quad (1)$$

where  $x_t = (1 - t)x + ty$ .

The above definition is inspired by the analogous upper sign-continuity for set-valued map, originated in [21], and given as follows

$T : X \rightrightarrows X^*$  is said to be upper sign-continuous at  $x \in \text{Dom } T$  if, for any  $v \in X$ , the following implication holds:

$$\left( \forall t \in ]0, 1[, \quad \inf_{x_t^* \in T(x_t)} \langle x_t^*, v \rangle \geq 0 \right) \Rightarrow \sup_{x^* \in T(x)} \langle x^*, v \rangle \geq 0$$

where  $x_t = x + tv$ .

A local version of the upper sign property of a bifunction has been considered in [13] by setting that  $f$  has the local upper sign property at  $x$  if (1) holds for every  $y$  in a certain neighbourhood  $B(x, \rho)$  of  $x$ , with  $\rho > 0$ . Clearly the upper sign property implies the local upper sign property. Actually, as shown in the forthcoming lemma, the upper sign property and its local counterpart coincide under mild assumptions.

**Lemma 2.2.** *Let  $K$  be a nonempty convex subset of  $X$  and  $f$  be a bifunction satisfying (H<sub>1</sub>) and the following property:*

$$\left\{ \begin{array}{l} \text{for every } x, y \in K, \text{ one has} \\ f(x, x) = 0 \text{ and } f(x, y) < 0 \Rightarrow f(x, z) < 0, \quad \forall z \in ]x, y[. \end{array} \right. \quad (2)$$

Then  $f$  has the local upper sign property on  $K$  if and only if  $f$  has the upper sign property on  $K$ .

**Proof.** Let us suppose that  $f$  has the locally upper sign property on  $K$  (with  $r > 0$ ) and let  $y \in K$  be such that  $f(x_t, x) \leq 0$ , for every  $x_t = tx + (1 - t)y$ ,  $t \in ]0, 1[$ . Then one immediately has that for every  $t \in ]0, 1[$ ,  $f(z_t, x) \leq 0$  where  $z_t = tx + (1 - t)z$  with  $z \in B(x, r) \cap K \cap ]x, y[$ . Now by the local upper sign property  $f(x, z) \geq 0$ . This implies that  $f(x, y) \geq 0$  because otherwise, in view of  $(H_1)$  and property (2), a contradiction occurs.  $\square$

**Remark 2.3.** As it can be easily shown, any bifunction satisfying assumption  $(H_2)$  also verifies property (2). This is in particular the case when  $f = f_T$  is a representative bifunction of a set-valued map  $T$ .

Now from the definition of  $f_T$  one clearly has the following equivalence for any set-valued map  $T : X \rightarrow 2^{X^*}$

$$T \text{ is upper sign-continuous at } x \Leftrightarrow \begin{cases} f_T \text{ has the upper} \\ \text{sign property at } x. \end{cases} \quad (3)$$

As a direct consequence of the classical Ky Fan intersection theorem [18] one can deduce an existence result for  $CFP(f, K)$ .

**Proposition 2.4.** *Let  $K$  be weakly compact and  $f$  be a properly quasimonotone equilibrium bifunction such that for every  $x \in K$  the subset  $\{y \in K : f(x, y) \leq 0\}$  is weakly closed. Then  $CFP(f, K)$  is nonempty.*

**Proof.** It follows the same lines as in [16, Th. 5.1]. Indeed, observe first that  $[\cap_{x \in K} F(x)] = CFP(f, K)$ , where, for every  $x \in K$ ,  $F(x)$  stands for the set  $F(x) = \{y \in K : f(x, y) \leq 0\}$ . On the other hand, the proper quasimonotonicity of  $f$  can be reformulated as

$$\forall x_1, \dots, x_n \in K, \forall x \in \text{co}\{x_1, \dots, x_n\}, \quad x \in \cup_{i=1}^n F(x_i),$$

which is classically expressed as  $F$  being a KKM map. The result follows now immediately by applying the Ky Fan intersection theorem.  $\square$

Let us end this section with an elementary remark on the set  $CFP(f, K)$ .

**Remark 2.5.** From the definitions of  $(H_2)$  and  $(H_3)$ , it follows immediately that if  $K$  is a closed convex subset of  $X$  and  $f : K \times K \rightarrow \mathbb{R}$  satisfies  $(H_2)$  and  $(H_3)$  then  $CFP(f, K)$  is convex and closed. Indeed, in this case, any sublevel set  $F(x) = \{y \in K : f(x, y) \leq 0\}$  is convex and closed, providing that  $x$  is an element of  $K$ .

### 3. Problem relationships

#### 3.1. Canonical inclusions between (QMEP) and (QEP)

The relations between the solution sets  $M(T, K)$  and  $S(T, K)$  respectively of the Minty and Stampacchia variational inequalities has been extensively studied in the literature (see e.g. [4] for a recent survey), in particular because they play an important role in the proofs of stability and existence results. Our aim in this short subsection is to specify these relations in the context of quasi-equilibrium problems.

**Proposition 3.1.** *Let  $f : X \times X \rightarrow \mathbb{R}$  be any bifunction and consider the following conditions:*

- i)  $f$  has the upper sign property on  $X$*
- ii)  $\text{QMEP}(f, K) \subseteq \text{QEP}(f, K)$  for all set-valued map  $K : X \rightarrow 2^X$  with convex values.*
- iii) for all  $y \in X$ ,  $\text{QMEP}(f, K_y) \subseteq \text{QEP}(f, K_y)$  with the constraint map  $K_y$  defined by  $K_y(x) = [x, y]$ .*

*Then  $i) \Rightarrow ii) \Rightarrow iii)$  and the three conditions are equivalent if  $f$  satisfies  $(H_1)$  and  $(H_2)$ .*

**Proof.**  $i) \Rightarrow ii)$  Immediate. Indeed let  $K : X \rightarrow 2^X$  be a set-valued map with convex values and take  $x \in \text{QMEP}(f, K)$ . For any  $y \in K(x)$  and any  $t \in ]0, 1[$  one clearly has  $f(tx + (1 - t)y, x) \leq 0$  and the conclusion follows by the upper sign property of  $f$ .

Assume now that  $iii)$  holds and suppose that  $f$  doesn't have the upper sign property on  $X$ . Therefore, one can find  $x, y \in X$  such that  $f(x, y) < 0$  and  $f(x_t, x) \leq 0$ , with  $x_t = tx + (1 - t)y$  for all  $t \in ]0, 1[$ . Hence, considering any element  $\tilde{y}$  in  $]x, y[$ , we get from  $(H_1)$  and (2) that  $f(x, \tilde{x}_t) < 0$ , with  $\tilde{x}_t = tx + (1 - t)\tilde{y}$  for all  $t \in ]0, 1[$ . On the other hand  $f(\tilde{x}_t, x) \leq 0$  for any  $t \in [0, 1]$  and therefore  $x \in \text{QMEP}(f, K_{\tilde{y}}(x))$  and thus, according to  $iii)$ ,  $x \in \text{QEP}(f, K_{\tilde{y}}(x))$ , implying that, for any  $t \in ]0, 1[$ ,  $f(x, \tilde{x}_t) \geq 0$ , which is a contradiction.  $\square$

As a direct consequence of Proposition 3.1, we obtain a sufficient condition for the upper-sign continuity of a set-valued map in terms of the inclusions between the solution sets of the Minty and Stampacchia defined by the set-valued map on intervals, thus providing a counterpart of [32, Th. 2] where these inclusions characterize the pseudomonotonicity of the set-valued map.

**Corollary 3.2.** *Let  $T : X \rightarrow 2^{X^*}$  be a set-valued map with convex compact values, and  $K$  be a convex subset of  $X$ . If  $M(T, [x, y]) \subseteq S(T, [x, y])$  for all  $x, y \in K$  then  $T$  is upper sign-continuous on  $K$ .*

**Proof.** Observing that, thanks to a classical Sion minimax theorem (see e.g. [31]), for all  $x, y \in K$ ,  $S(T, [x, y]) = \text{EP}(f_T, [x, y])$  and  $M(T, [x, y]) =$

$MEP(f_T, [x, y])$ , one can deduce that  $MEP(f_T, [x, y]) \subseteq EP(f_T, [x, y])$ . Therefore, by Proposition 3.1,  $f_T$  has the upper sign property on  $K$  and thus, according to (3),  $T$  is upper sign-continuous on  $K$ .  $\square$

### 3.2. Variational formulation of equilibrium problems

As quoted previously, any Stampacchia variational inequality can be reformulated as an equilibrium problem, thanks to the use of the representative bifunction  $f_T$ . This subsection is devoted to the study of the reverse transformation and more precisely, given a bifunction  $f$  and a subset  $K$ , to give sufficient conditions under which the solution set  $EP(f, K)$  of the equilibrium problem coincides with the solution set of a certain Stampacchia variational inequality.

Associated to a convex subset  $K$  of  $X$  and a bifunction  $f : X \times X \rightarrow \mathbb{R}$ , we define the set-valued map  $N : K \rightarrow 2^{X^*}$  as

$$N(x) = \{x^* \in X^* : \langle x^*, y - x \rangle \leq 0, \forall y \in L_f(x)\},$$

where  $L_f(x)$  stands for the sublevel set  $L_f(x) = \{z \in K : f(x, z) \leq 0\}$ .

**Proposition 3.3.** *Let  $K$  be a convex subset of  $X$  and  $f : X \times X \rightarrow \mathbb{R}$  be a quasimonotone bifunction. Then the set-valued map  $N$  is quasimonotone on  $K$ .*

**Proof.** Let  $x$  and  $y$  be two elements of  $K$  and  $x^* \in N(x)$  be such that  $\langle x^*, y - x \rangle > 0$ . From the definition of  $N$  one gets that  $f(x, y) > 0$ , and hence  $f(y, x) \leq 0$  by quasimonotonicity of  $f$ . Now since  $x$  is an element of  $L_f(y)$ , one has  $\langle y^*, x - y \rangle \leq 0$  for all  $y^* \in N(y)$ , proving that  $N$  is quasimonotone.  $\square$

Following a technique used in [7], let us define now a normalized version of the set-valued map  $N$  by  $D : K \rightarrow 2^{X^*}$ :

$$D(x) = \begin{cases} \text{co}(N(x) \cap S^*) & \text{if } x \notin \arg \min_X f(x, \cdot) \\ B^* & \text{otherwise.} \end{cases}$$

By the same proof as in [5, Lemma 3.1], one can prove that if  $f$  satisfies  $(H_1)$  and  $(H_2)$  then one has  $0 \notin D(x)$ , for all  $x \notin \arg \min_X f(x, \cdot)$ .

**Proposition 3.4.** *Let  $K$  be a convex subset of  $X$  and  $f : X \times X \rightarrow \mathbb{R}$  be a quasimonotone bifunction satisfying  $(H_1)$ ,  $(H_2)$  and  $(H_3)$ . If  $\text{int}(L_f(x)) \neq \emptyset$  for any  $x \notin \arg \min_X f(x, \cdot)$ , then  $EP(f, K) = S(D, K)$ .*

**Proof.** If  $x \in EP(f, K)$  then  $x$  is actually a minimizer of  $f(x, \cdot)$  on  $K$ . Now, if  $x \in \arg \min_X f(x, \cdot)$  then  $0 \in D(x)$ , i.e.,  $x \in S(D, K)$ . On the other hand, if  $x \notin \arg \min_X f(x, \cdot)$  then, in view of the first order necessary optimality condition in quasiconvex programming (see Theorem 4.1 in [8]), there exists

$x^* \in D(x)$  such that  $x$  is a solution of  $S(D, K)$ .

Conversely, assume that  $x \in S(D, K)$ . If  $0 \in D(x)$  then  $x$  belongs to  $\arg \min_X f(x, \cdot)$ , so that  $f(x, y) \geq 0$  for all  $y \in K$ , i.e.,  $x \in EP(f, K)$ . Now if  $0 \notin D(x)$  then there exists a non zero element  $x^*$  of  $N(x)$  such that  $\langle x^*, y - x \rangle \geq 0$ , for every  $y \in K$  and thus, according to the sufficient optimality condition given in [7, Corollary 4.5], one gets that  $x$  is a minimizer of  $f(x, \cdot)$  and thus  $x \in EP(f, K)$ .  $\square$

#### 4. Existence of quasi-equilibriums

Contrary to the proofs used for quasimonotone variational inequalities in [6], for quasimonotone equilibrium problems in [13] and for quasi-equilibrium problems in [22] which are based on KKM techniques, the existence result proposed in the forthcoming Theorem 4.5 is proved by mixing a Kakutani fixed point theorem with an adapted study of the stability (regularity) of the solution set (map) of a perturbed Convex Feasible Problem.

So let us first consider the following perturbation of the Convex Feasibility Problem: let  $K : \mathbb{R}^m \rightarrow 2^{\mathbb{R}^n}$  be a set-valued map and  $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  a bifunction. Then for any  $\mu \in \mathbb{R}^m$ , we define

$$(CFP_\mu) \quad \text{find } x \in K(\mu) \text{ such that } f(y, x) \leq 0 \quad \forall y \in K(\mu).$$

Let us denote by  $CFP(f, K(\cdot))$  the solution map which associates to any  $\mu$  the solution set  $CFP(f, K(\mu))$  of problem  $(CFP_\mu)$ . The following proposition, which is an adaptation of Proposition 4.4 in [4], gives sufficient conditions for the set-valued map  $CFP(f, K(\cdot))$  to have a closed graph.

**Proposition 4.1.** *Let  $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a bifunction and  $K : \mathbb{R}^m \rightarrow 2^{\mathbb{R}^n}$  be a set-valued map which satisfy the following properties*

- i)  $f$  verifies  $(H_3)$ ;
- ii) for all sequence  $(y_k)_k \subset \mathbb{R}^n$  converging to  $y \in \mathbb{R}^n$  and such that  $\liminf_{k \rightarrow +\infty} f(y_k, x) \leq 0$ , one has  $f(y, x) \leq 0$ ;
- iii)  $K$  is closed, lower semicontinuous and convex valued, with  $\text{int}(K(\mu)) \neq \emptyset$ , for all  $\mu \in \text{dom } K$ .

*Then the solution map  $CFP(f, K(\cdot))$  is closed.*

We mention that the assumptions on the map  $K$  in item iii) above are rather mild and quite natural. For instance, they hold when  $K(\mu)$  is of the form  $K(\mu) = \{x \in \mathbb{R}^n : h(x) \leq \mu\}$ , where  $h : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is continuous.

In order to prove the above proposition let us recall the following interesting properties, established in [1] and [4].

**Proposition 4.2 ([1, Prop. 3.2]).** *Let  $S$  be a subset of  $X$  and  $(S_k)_k$  be a sequence of convex subsets of  $X$ . Assume that  $\text{int}(S_k) \neq \emptyset$  for all  $k$  and that  $\text{int}(S) \neq \emptyset$ . Then  $i) \Rightarrow ii)$  where*

- i)  $w\text{-}\limsup_{k \rightarrow +\infty} S_k \subset S \subset \liminf_{k \rightarrow +\infty} S_k$
- ii)  $w\text{-}\limsup_{k \rightarrow +\infty} S_k \subset S$ ,  $\text{int}(S) \subset \liminf_{k \rightarrow +\infty} \text{int}(S_k)$  and  $S$  is convex.

**Proposition 4.3** ([4, Lemma 4.1]). *Let  $C$  and  $(C_k)_k$  be a convex set and a sequence of convex subsets of  $\mathbb{R}^n$  such that  $C \subset \liminf_{k \rightarrow +\infty} \text{int}(C_k)$ . If  $\text{int}(C) \neq \emptyset$  and  $\text{int}(C_k) \neq \emptyset$ , for all  $k \in \mathbb{N}$ , then for each  $y \in \text{int}(C)$  there exists  $k_0 \in \mathbb{N}$  such that  $y \in \text{int}(C_k)$ , for all  $k \geq k_0$ .*

**Proof of Proposition 4.1.** Let  $((\mu_k, x_k))_k \subset \text{Gr } CFP(f, K(\cdot))$  be a sequence converging to  $(\mu, x)$ . Since  $K$  is closed, one has that  $x \in K(\mu)$ .

We claim that any  $y \in \text{int}(K(\mu))$  belongs to  $\text{int}(K(\mu_k))$  for  $k$  large enough. Observe that, according to the lower semicontinuity of  $K$ ,  $K(\mu) \subset \liminf_{k \rightarrow +\infty} K(\mu_k)$ . Since the sets  $K(\mu)$  and  $K(\mu_k)$  are convex with nonempty interior, it follows from Proposition 4.2 that

$$y \in \text{int}(K(\mu)) \subset K(\mu) \subset \liminf_{k \rightarrow +\infty} \text{int}(K(\mu_k)),$$

and thus the claim is now a consequence of Proposition 4.3.

Since, for all  $k$  we have  $x_k \in CFP(f, K(\mu_k))$ , it follows that

$$f(y, x_k) \leq 0,$$

and therefore  $f(y, x) \leq 0$ , by property  $(H_3)$ . Now, for any  $y \in K(\mu)$  there exists a sequence  $(y_k)_k \subset \text{int}(K(\mu))$  converging to  $y$ . So

$$f(y_k, x) \leq 0, \text{ for any } k \text{ large enough.}$$

By assumption ii) on the bifunction  $f$ , one has that  $f(y, x) \leq 0$  and therefore  $x \in CFP(f, K(\mu))$ .  $\square$

**Remark 4.4.** An analogous to property ii) of the bifunction  $f$  in Proposition 4.1 was introduced in [1, 4] for set-valued mappings, and it was called *dually lower semicontinuity* in [4].

We are now in a position to establish our existence result for properly quasi-monotone quasi-equilibrium problems.

**Theorem 4.5.** *Let  $f : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a bifunction,  $C$  a nonempty convex compact subset of  $\mathbb{R}^n$  and  $K : C \rightarrow 2^C$  be a set-valued map, and suppose that the following properties hold:*

- i) *the map  $K$  is closed and lower semicontinuous, with convex values, and  $\text{int}(K(x)) \neq \emptyset$  for all  $x \in C$ ,*
- ii)  *$f$  has the upper sign property on  $C$ ,*
- iii) *for all  $x, y \in \mathbb{R}^n$  and all sequence  $(y_k)_k \subset \mathbb{R}^n$  converging to  $y$  it holds that*

$$\liminf_{k \rightarrow +\infty} f(y_k, x) \leq 0 \Rightarrow f(y, x) \leq 0,$$

iv)  $f$  is properly quasimonotone and satisfies  $(H_2)$  and  $(H_3)$ .

Then the quasi-equilibrium problem  $QEP(f, K)$  admits at least a solution.

**Proof.** Since  $f$  is properly quasimonotone and  $K$  is nonempty, convex and compact valued, it follows from Proposition 2.4 that  $CFP(f, K(x)) \neq \emptyset$  for all  $x \in C$ . Now the map  $CFP(f, K(\cdot)) : C \rightarrow 2^C$  is closed and has convex and closed values by Proposition 4.1 and Remark 2.5. Hence, due to the compactness of  $C$ , this solution map is upper semicontinuous. Now according to the Kakutani Fixed Point Theorem (see e.g. [28]), we conclude that the map  $CFP(f, K(\cdot))$  admits at least a fixed point  $x$ , or in other words  $x \in CFP(f, K(x))$ . By Proposition 3.1 *ii*) we get that  $x \in EP(f, K(x))$ , and therefore  $x$  is a solution of  $QEP(f, K)$ .  $\square$

**Remark 4.6.** i) Note that the above result is intrinsically limited to the finite dimensional case because the values of the constraint map  $K$  are assumed to be compact and with nonempty interior.

ii) Since, in the above theorem,  $f$  is assumed to be properly quasimonotone and to have the upper sign property, property  $(H_1)$  immediately holds.

iii) The upper sign property and hypothesis *iii*) in Theorem 4.5 are independent. Indeed, the bifunction  $f_1 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ , defined by  $f_1(x, y) = \min\{x, y\}$ , is continuous and hence it satisfies hypothesis *iii*) in Theorem 4.5, but  $f_1$  does not enjoy the upper sign property. Conversely, the bifunction  $f_2 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ , defined by  $f_2(x, y) = \begin{cases} x-y & \text{if } x \neq y \\ 1 & \text{if } x = y \end{cases}$ , has the upper sign property on  $\mathbb{R}$ . However,  $f_2$  does not satisfy condition *iii*) in Theorem 4.5.

**Acknowledgements.** The authors would like to thanks the referee for valuable remarks and comments which improved the quality of the paper.

## References

- [1] M. Ait Mansour, D. Aussel: Quasimonotone variational inequalities and quasi-convex programming: qualitative stability, *J. Convex Analysis* 15 (2008) 459–472.
- [2] D. Aussel, R. Correa, M. Marechal: Gap functions for quasivariational inequalities and generalized Nash equilibrium problems, *J. Optim. Theory Appl.* 151 (2011) 474–488.
- [3] D. Aussel, J. Cotrina: Quasimonotone quasivariational inequalities: existence results and applications, *J. Optim. Theory Appl.* 158 (2013) 637–652.
- [4] D. Aussel, J. Cotrina: Stability of quasimonotone variational inequalities under sign-continuity, *J. Optim. Theory Appl.* 158 (2013) 653–667.
- [5] D. Aussel, J. Dutta: On gap functions for multivalued Stampacchia variational inequalities, *J. Optim. Theory Appl.* 149 (2011) 513–527.

- [6] D. Aussel, N. Hadjisavvas: On quasimonotone variational inequalities, *J. Optim. Theory Appl.* 121 (2004) 445–450.
- [7] D. Aussel, N. Hadjisavvas: Adjusted sublevel sets, normal operator and quasi-convex programming, *SIAM J. Optim.* 16 (2005) 358–367.
- [8] D. Aussel, J. J. Ye: Quasiconvex programming with locally starshaped constraint region and applications to quasiconvex MPEC, *Optimization* 55 (2006) 433–457.
- [9] M. Bianchi, R. Pini: A note on equilibrium problems with properly quasimonotone bifunctions, *J. Glob. Optim.* 20 (2001) 67–76.
- [10] M. Bianchi, R. Pini: Coercivity conditions for equilibrium problems, *J. Optim. Theory Appl.* 124 (2005) 79–92.
- [11] M. Castellani, M. Giuli: On equivalent equilibrium problems, *J. Optim. Theory Appl.* 147 (2010) 157–168.
- [12] M. Castellani, M. Giuli: Pseudomonotone diagonal subdifferential operators, *J. Convex Analysis* 20 (2013) 1–12.
- [13] M. Castellani, M. Giuli: Refinements of existence results for relaxed quasimonotone equilibrium problems, *J. Glob. Optim.* 57 (2013) 1213–1227.
- [14] M. Castellani, M. Giuli: An existence result for quasiequilibrium problems in separable Banach spaces, *J. Math. Anal. Appl.* 425 (2015) 85–95.
- [15] J. P. Crouzeix, E. Ocaña Anaya: Maximality is nothing but continuity, *J. Convex Analysis* 17 (2010) 521–534.
- [16] A. Daniilidis, N. Hadjisavvas: Characterization of nonsmooth semistrictly quasiconvex and strictly quasiconvex functions, *J. Optim. Theory Appl.* 102 (1999) 525–536.
- [17] F. Facchinei, J. S. Pang: *Finite Dimensional Variational Inequalities*, Springer, New York (2003).
- [18] K. Fan: A generalization of Tychonoff’s fixed point theorem, *Math. Ann.* 142 (1961) 305–310.
- [19] F. Flores-Bazán: Existence theorems for generalized noncoercive equilibrium problems: The quasiconvex case, *SIAM J. Optim.* 11 (2000) 675–790.
- [20] F. Flores-Bazán: Existence theory for finite dimensional pseudomonotone equilibrium problems, *Acta Appl. Math.* 77 (2003) 249–297.
- [21] N. Hadjisavvas: Continuity and maximality properties of pseudomonotone operators, *J. Convex Analysis* 10 (2003) 459–469.
- [22] N. X. Hai, P. Q. Khanh: Existence of solutions to general quasiequilibrium problems and applications, *J. Optim. Theory Appl.* 133 (2007) 317–327.
- [23] A. Iusem: On the maximal monotonicity of diagonal subdifferential operators, *J. Convex Analysis* 18 (2011) 489–503.
- [24] A. Iusem, G. Kassay, W. Sosa: On certain conditions for the existence of solutions of equilibrium problems, *Math. Program.* 116 (2009) 259–273.
- [25] A. Iusem, G. Kassay, W. Sosa: An existence result for equilibrium with some surjectivity consequences, *J. Convex Analysis* 16 (2009) 807–826.

- [26] A. Iusem, W. Sosa: New existence results for equilibrium problems, *Nonlinear Anal.* 52 (2003) 621–635.
- [27] A. Iusem, B. F. Svaiter: On diagonal subdifferential operators in nonreflexive Banach spaces, *Set-Valued Var. Anal.* 20 (2012) 1–14.
- [28] S. Kakutani: A generalization of Brouwer’s fixed-point theorem, *Duke Math. J.* 8 (1941) 457–459.
- [29] L. J. Lin, M. F. Yang, Q. H. Ansari, G. Kassay: Existence results for Stampacchia and Minty type implicit variational inequalities with multivalued maps, *Nonlinear Anal.* 61 (2005) 1–19.
- [30] J. S. Pang, M. Fukushima: Quasi-variational inequalities, generalized Nash equilibria and multi-leader-follower games, *Comput. Manag. Sci.* 2 (2005) 21–56.
- [31] M. Sion: On general minimax theorems, *Pac. J. Math.* 8 (1958) 171–176.
- [32] J. Reinhard: A note on Minty variational inequalities and generalized monotonicity, in: *Generalized Convexity and Generalized Monotonicity* (Samos, 1999), N. Hadjisavvas et al. (ed.), *Lecture Notes in Economics and Mathematical Systems* 502, Springer, Berlin (2001) 240–246.