

Existence and Stability for Generalized Polynomial Vector Variational Inequalities

Tran Van Nghi

*Hanoi Pedagogical University 2, Hanoi, Vietnam
tranvanngghi@hpu2.edu.vn*

Nguyen Nang Tam*

*Institute of Theoretical and Applied Research, Duy Tan University, Hanoi, Vietnam;
and: Faculty of Natural Sciences, Duy Tan University, Da Nang, Vietnam
nguyennangtam@duytan.edu.vn*

Received: June 30, 2022

Accepted: July 10, 2023

We investigate the generalized polynomial vector variational inequality (GPVVI), which is a natural generalization of generalized polynomial variational inequality (GPVI) and vector variational inequality (VVI). Due to the scalarization method, which is a powerful technique in vector optimization, we establish a relationship between the Pareto solution sets of the GPVVI and the solution set of the GPVI. By using the concept on exceptional family of elements, recession cone, and positive semi-definiteness of matrices, we present sufficient conditions for the nonemptiness and boundedness of the Pareto solution sets of the GPVVI. We present sufficient conditions for the upper/lower semicontinuity of the weak Pareto solution map and the stability for GPVVI. Finally, we obtain some applications to polynomial variational inequality. The presented result develops and complements the previous ones.

Keywords: General vector variational inequality, Pareto solution, solution existence, stability, upper/lower semicontinuity.

2020 Mathematics Subject Classification: 90C29, 90C31, 90C33, 54C60, 49J40, 49J53, 49K40.

1. Introduction

Variational inequalities (VI) theory, which was introduced by Stampacchia [25], is an important branch of the mathematical sciences. The concept of *vector variational inequality* (VVI), which is an adequate tool for studying vector optimization problems and vector equilibrium problems, was proposed by Giannessi [3]. By using a stability theorem of Robinson and a scalarization method, Yen and Yao [31] established sufficient conditions for the upper semicontinuity of the solution maps of parametric monotone affine vector variational inequalities (AVVI). A survey of recent results on the VVI was given in [4]. Yen [30] showed that the importance of VVIs for vector optimization is the same as that of variational inequalities for scalar optimization.

The *general variational inequality* (GVI), which was firstly proposed by Noor [21], has received considerable attention in recent three decades. Noor [22] also showed that the minimum of a differentiable *hg*-convex function on the *hg*-convex set K

* Corresponding author.

in $\mathbb{R}^n$ can be characterized by the GVI. It is well-known that the problem GVI is equivalent to a class of the fixed point problems. These equivalent formulations allow us to approach studying polynomial optimization and fixed point theory via the GVI. Moreover, the GVI contains the class of inverse variational inequalities (see [13, 14]). Recently, existence and stability for generalized polynomial variational inequalities (GPVI), GVI, VI, and general polynomial complementarity problems have been studied (see [12, 15–20, 27, 29]). The obtained results have extended ones in [23].

In this paper, we study the *generalized polynomial vector variational inequality* (GPVVI), which is a natural generalization of the GPVI and the VVI. The major contributions to the GPVVI theory are listed as follows:

- (i) A relationship between the Pareto solution sets of the GPVVI and the solution set of the GPVI is established by using the scalarization method, which is a powerful technique in vector optimization.
- (ii) By using the concept on exceptional family of elements, recession cone, and positive semi-definiteness of matrices, we present sufficient conditions for the nonemptiness and boundedness of the Pareto solution sets of the GPVVI;
- (iii) We propose sufficient conditions for the upper/lower semicontinuity of the weak Pareto solution map and characterize the stability for GPVVI;
- (iv) Finally, we give some applications to polynomial variational inequality.

Our result develops and complements the previous ones for VIs, GPVIs and VVIs in [8, 11, 15, 26, 28, 31].

The outline of the paper is as follows. Section 2 gives some preliminaries. The existence of Pareto solutions is established in Section 3. Due to this result, we present sufficient conditions for upper/lower continuity of the weak Pareto solution map of the GPVVI in Section 4. Some stability results for the GPVVI are also proposed Section 4. Some applications to polynomial variational inequality are given in the last section.

2. Preliminaries

Let $\mathbb{R}^s$ be s -dimensional Euclidean space equipped with the standard scalar product and the Euclidean norm, $\mathbb{R}_S^{s \times s}$ be the space of real symmetric $(s \times s)$ -matrices equipped with the matrix norm induced by the vector norm in $\mathbb{R}^s$. The scalar product of vectors x, y and the Euclidean norm of a vector x in a finite-dimensional Euclidean space are denoted, respectively, by $\langle x, y \rangle$ and $\|x\|$. Vectors in Euclidean spaces are interpreted as columns of real numbers. The notation $x \geq y$ (resp., $x > y$) means that every component of x is greater or equal (resp., greater) the corresponding component of y . For each natural number q , denote $[q] := \{1, \dots, q\}$. The norm in the product space $X_1 \times \dots \times X_k$ of the normed spaces $X_1, \dots, X_k$ is set to be

$$\|(x_1, \dots, x_k)\| = (\|x_1\|^2 + \dots + \|x_k\|^2)^{\frac{1}{2}}.$$

Let K be a nonempty closed convex set of $\mathbb{R}^s$, and $F_i, G : \mathbb{R}^s \rightarrow \mathbb{R}^s$, $i = 1, \dots, m$, be polynomial mappings. Let $F_i = (F_{i1}, \dots, F_{is})$, $i = 1, \dots, m$, $G = (G_1, \dots, G_s)$, where for each $j = 1, \dots, s$, the j -th component F_{ij} of F_i and G_j of G are polynomial functions in the variables $x_1, \dots, x_s$.

Let $F = (F_1, \dots, F_m)$. For each $x \in K$ and $h \in \mathbb{R}^s$, denote

$$F(x)(h) := (\langle F_1(x), h \rangle, \dots, \langle F_m(x), h \rangle).$$

The so-called *generalized polynomial vector variational inequality* (GPVVI) defined by (F, G, K) is to find a vector $\bar{x} \in \mathbb{R}^s$ such that

$$G(\bar{x}) \in K \text{ and } F(\bar{x})(x - G(\bar{x})) \notin -\mathbb{R}_+^m \setminus \{0\} \quad \forall x \in K, \quad (\text{GPVVI}(F, G, K))$$

where $\mathbb{R}_+^m$ is the nonnegative orthant of $\mathbb{R}^m$. To this problem one associates the following: Find a vector $\bar{x} \in \mathbb{R}^s$ such that

$$G(\bar{x}) \in K \text{ and } F(\bar{x})(x - G(\bar{x})) \notin -\text{int}(\mathbb{R}_+^m) \quad \forall x \in K, \quad (\text{GPVVI}^w(F, G, K))$$

where $\text{int}(\mathbb{R}_+^m)$ is the topological interior of $\mathbb{R}_+^m$.

The solution set of $\text{GPVVI}(F, G, K)$ and $\text{GPVVI}^w(F, G, K)$ are denoted, respectively, by $Sol(F, G, K)$ and $Sol^w(F, G, K)$. The elements of the first set (resp., the second set) are said to be the *Pareto solutions* (resp., the *weak Pareto solutions*) of $\text{GPVVI}(F, G, K)$.

For the case where $G = id_{\mathbb{R}^s}$, the problem $\text{GPVVI}(F, G, K)$ reduces to the following *polynomial vector variational inequality* (VVI) (see [9])

$$\text{Find } \bar{x} \in K \text{ s.t. } F(\bar{x})(x - \bar{x}) \notin -\mathbb{R}_+^m \setminus \{0\} \quad \forall x \in K. \quad (\text{VVI}(F, K))$$

If $m = 1$ and $F = F_1$ then the problem $\text{GPVVI}(F, G, K)$ reduces to the following *generalized polynomial variational inequality* (GPVI)

$$\text{Find } \bar{x} \in K \text{ s.t. } \langle F(\bar{x}), x - G(\bar{x}) \rangle \geq 0 \quad \forall x \in K. \quad (\text{GVI}(F, G, K))$$

which was first proposed by Noor [21]. In particular, if $G = id_{\mathbb{R}^s}$, $m = 1$, and $F = F_1$ then $\text{GPVVI}(F, G, K)$ reduces to the following *polynomial variational inequality* (PVI) (see [5, 10])

$$\text{Find } \bar{x} \in K \text{ s.t. } \langle F(\bar{x}), x - \bar{x} \rangle \geq 0 \quad \forall x \in K. \quad (\text{PVI}(F, K))$$

For any nonempty closed convex set K of $\mathbb{R}^s$, the asymptotic (recession) cone of K is denoted by

$$K^\infty = \{v \in \mathbb{R}^s : x + tv \in K \quad \forall t \geq 0\}.$$

For any cone $C \subset \mathbb{R}^s$, the dual of C is denoted by

$$C^* := \{y \in \mathbb{R}^s : \langle h, y \rangle \geq 0 \quad \forall h \in C\}.$$

For any given polynomial p , the *degree of p* is denoted by $\deg p$. Let $P : \mathbb{R}^s \rightarrow \mathbb{R}^s$ be a polynomial mapping, i.e. $P = (p_1, \dots, p_s)$, where p_i is the polynomial in s variables, $i = 1, \dots, s$. The maximum of the numbers $\deg p_i$ is called the degree of the polynomial map P and denoted by $\deg P$. For any polynomial mapping $P = (p_1, \dots, p_s)$ with $\deg P = d$, we denote $P^\infty = (p_1^\infty, \dots, p_s^\infty)$, where p_i^∞ is the homogeneous component of degree d of p_i , $i = 1, \dots, s$, and

$$\text{Ker}(P^\infty) = \{x \in \mathbb{R}^s : P^\infty(x) = 0\}.$$

Note that the polynomial mapping P^∞ is homogeneous of degree d and

$$P^\infty(x) = \lim_{t \rightarrow +\infty} \frac{P(tx)}{t^d} \quad \forall x \in \mathbb{R}^s.$$

The polynomial mapping P^∞ is called the *leading term* of the polynomial mapping P .

Let $\mathcal{P}^{[d,s]}$ be the linear space of all polynomial mappings $P = (p_1, \dots, p_s)$ of degree at most d from $\mathbb{R}^s$ to $\mathbb{R}^s$ with the norm defined by

$$\|P\| = (\|p_1\|^2 + \dots + \|p_s\|^2)^{\frac{1}{2}}$$

for all $P = (p_1, \dots, p_s) \in \mathcal{P}^{[d,s]}$, where $\|p_i\|$ denotes the norm of polynomial p_i . It is clear that, if $\{P^k\}$ is a convergent sequence in $\mathcal{P}^{[d,s]}$ with $P^k \rightarrow P$ and $\{x^k\}$ is a convergent sequence in $\mathbb{R}^s$ with $x^k \rightarrow x$, then $P^k(x^k) \rightarrow P(x)$ and $(P^k)^\infty \rightarrow P^\infty$.

Let $\deg F_{ij} = \delta^{ij}$, $\deg G_j = \sigma^j$. Let $\deg F_i = \delta^i := \max\{\delta^{ij} : j = 1, \dots, s\}$, $\deg G = \sigma := \max\{\sigma^j : j = 1, \dots, s\}$, define $\deg F = \delta := \max\{\delta^i : i = 1 \dots m\}$.

Let

$$\Omega := \underbrace{\mathcal{P}^{[\delta,s]} \times \dots \times \mathcal{P}^{[\delta,s]}}_{m \text{ times}} \times \mathcal{P}^{[\sigma,s]}$$

A multifunction $\Phi : \Omega \rightrightarrows \mathbb{R}^s$ is said to be *upper semicontinuous* at $\bar{z} \in \Omega$ if for each open set V containing $\Phi(\bar{z})$ there exists $\delta > 0$ such that $\Phi(z) \subset V$ for every $z \in \Omega$ satisfying $\|z - \bar{z}\| < \delta$. We also say that Φ is *lower semicontinuous* at $\bar{z}$ if $\Phi(\bar{z}) \neq \emptyset$ and, for each open set V satisfying $\Phi(\bar{z}) \cap V \neq \emptyset$, there exists $\epsilon > 0$ such that $\Phi(z) \cap V \neq \emptyset$ for every $z \in \Omega$ satisfying $\|z - \bar{z}\| < \epsilon$. If Φ is lower and upper semicontinuous at $\bar{z}$ then Φ is called *continuous* at $\bar{z}$.

Let F, G be two continuous functions. A set of points $\{x^k\} \subset \mathbb{R}^s$ is called an *exceptional family of elements* (see [29]) for the pair (F, G) with respect to $\bar{x} \in \mathbb{R}^s$ if $\|x^k\| \rightarrow \infty$ as $k \rightarrow \infty$; and for each x^k , there exists a scalar $\alpha^k > 0$ such that $z^k := \alpha^k(x^k - \bar{x}) + G(x^k) \in K$ and

$$-\alpha^k(x^k - \bar{x}) - F(x^k) \in \mathcal{N}_K(z^k),$$

where $\mathcal{N}_K(z^k)$ is the normal cone of K at z^k .

For any nonempty closed convex set K of $\mathbb{R}^s$, the *projection* of $\hat{x}$ on K is denoted by $\Pi_K(\hat{x})$.

For any continuous mapping $g : \mathbb{R}^s \rightarrow \mathbb{R}^s$ and $\Omega_r^{\hat{x}} := \{x \in \mathbb{R}^s : \|g(x)\| < r\}$, where $r > \|\Pi_K(\hat{x})\|$ for any given $\hat{x} \in \mathbb{R}^s$, the *topological degree of g at $\Pi_K(\hat{x})$ relative to $\Omega_r^{\hat{x}}$* is denoted by $\deg(g(\cdot), \Omega_r^{\hat{x}}, \Pi_K(\hat{x}))$.

Let K be a nonempty closed convex set of $\mathbb{R}^s$ and $g : \mathbb{R}^s \rightarrow \mathbb{R}^s$ be a continuous mapping. The pair (g, K) is said to have the *property $\mathcal{A}$* if it satisfies the following conditions:

- (a₁) the boundedness of $\|g(x)\|$ implies the boundedness of $\|x\|$;
- (a₂) $\deg(g(\cdot), \Omega_r^{\hat{x}}, \Pi_K(\hat{x}))$ is defined and nonzero.

We have the following lemma.

Lemma 2.1. *Consider the problem $\text{GVI}(f, g, K)$ with K being a nonempty closed convex set and $f, g : \mathbb{R}^s \rightarrow \mathbb{R}^s$ being continuous mappings. If the pair (g, K) has property $\mathcal{A}$, then there exists either a solution of $\text{GVI}(f, g, K)$ or an exceptional family of elements for the pair (f, g) with respects to any given $\hat{x} \in \mathbb{R}^s$.*

Proof. Its proof is similar to that of [7, Theorem 2.2] and [1, Theorem 1]. $\square$

We call f *copositive* (*strictly copositive*) with respect to g on K if

$$\langle f(x), g(x) \rangle \geq 0 \ (> 0)$$

for every $x \in \mathbb{R}^n$ satisfying $g(x) \in K$. The map f is called *copositive* (*strictly copositive*) on K if

$$\langle f(x), x \rangle \geq 0 \ (> 0) \ \forall x \in K.$$

We say that f is *strictly monotone* with respect to g on K if

$$\langle f(x) - f(y), g(x) - g(y) \rangle > 0$$

for every $x, y \in \mathbb{R}^n$ satisfying $x \neq y$ and $g(x), g(y) \in K$.

3. Existence of Pareto solutions

Our goal in this section is twofold. Firstly, we use the scalarization method to establish a relationship between the Pareto solution sets of the GCVI and the solution set of the GVI. Secondly, by the above relation and the exceptional family of elements, we present sufficient conditions for the nonemptiness and boundedness of the Pareto solution sets of the GPVVI. To do this, let

$$\Delta := \left\{ \xi = (\xi_1, \dots, \xi_m) \in \mathbb{R}_+^m : \sum_{i=1}^m \xi_i = 1 \right\}$$

and $ri\Delta := \{ \xi = (\xi_1, \dots, \xi_m) \in \Delta : \xi_i > 0 \ \forall i \in [m] \}$.

For some $\xi = (\xi_1, \dots, \xi_m) \in \Delta$, consider the following variational inequality

$$\text{Find } \bar{x} \in K \text{ s.t. } \left\langle \sum_{i=1}^m \xi_i F_i(\bar{x}), x - G(\bar{x}) \right\rangle \geq 0 \ \forall x \in K. \quad (\text{GVI}_\xi(F, G, K))$$

Denote by $\text{Sol}(\text{GVI}_\xi(F, G, K))$ the solution set of $(\text{GVI})_\xi(F, G, K)$, $F_\xi := \sum_{i=1}^m \xi_i F_i$. Denote by $F^\infty := (F_1^\infty, \dots, F_m^\infty)$ the leading term of F with degree δ , that is, $F^\infty(\lambda h) = \lambda^\delta F^\infty(h)$.

For given $\text{GPVVI}(F, G, K)$, $\text{GPVVI}^w(F, G, K)$, $\xi \in \Delta$ and $\rho \geq 0$ the solution set of $\text{GPVVI}(F_\xi^\infty + \rho G^\infty, G^\infty, K^\infty)$ and $\text{GPVVI}^w(F^\infty, G^\infty, K^\infty)$ are denoted, respectively, by $\text{Sol}(F_\xi^\infty + \rho G^\infty, G^\infty, K^\infty)$ and $\text{Sol}^w(F^\infty, G^\infty, K^\infty)$.

The following lemma shows a relationship between the Pareto solution sets of the inequality $\text{GPVVI}(F, G, K)$ and the solution set of $(\text{GVI})_\xi(F, G, K)$.

Lemma 3.1. *Let K be a nonempty closed convex set. It holds that*

$$\bigcup_{\xi \in ri\Delta} \text{Sol}(\text{GVI}_\xi(F, G, K)) \subset \text{Sol}(F, G, K) \quad (1)$$

$$\text{and} \quad \text{Sol}^w(F, G, K) = \bigcup_{\xi \in \Delta} \text{Sol}(\text{GVI}_\xi(F, G, K)). \quad (2)$$

Proof. To prove inclusion (1), we take any $\xi \in ri\Delta$ and $\bar{x} \in \text{Sol}(\text{GVI}_\xi(F, G, K))$. For every $x \in K$, we have $G(\bar{x}) \in K$ and

$$0 \leq \left\langle \sum_{i=1}^m \xi_i F_i(\bar{x}), x - G(\bar{x}) \right\rangle = \sum_{i=1}^m \xi_i \langle F_i(\bar{x}), x - G(\bar{x}) \rangle = \xi^T F_\xi(\bar{x})(x - \bar{x}). \quad (3)$$

Hence, there no exist $x \in K$ such that $G(\bar{x}) \in K$ and

$$F(\bar{x})(x - G(\bar{x})) \in -\mathbb{R}_+^m \setminus \{0\},$$

that is, $\bar{x} \in \text{Sol}(F, G, K)$.

We now show that (2) holds. Indeed, for any $\bar{x} \in \text{Sol}(\text{GVI}_\xi(F, G, K))$ with $\xi \in \Delta$, we have (3). Then, there no exist $\bar{x} \in \mathbb{R}^n$ satisfying $G(\bar{x}) \in K$ and $x \in K$ such that $F_\xi(\bar{x})(x - G(\bar{x})) \in -\text{int}(\mathbb{R}_+^m) \setminus \{0\}$. It implies $\bar{x} \in \text{Sol}^w(F, G, K)$. Conversely, for any $\bar{x} \in \text{Sol}^w(F, G, K)$, we have

$$\{F_\xi(\bar{x})(x - G(\bar{x})) : x \in K\} \cap -\text{int}(\mathbb{R}_+^s) = \emptyset$$

and $G(\bar{x}) \in K$. According to the separation theorem (see, for instance, [24, Theorem 11.3]), there exists $\hat{\xi} \in \mathbb{R}$ with $\|\hat{\xi}\| = 1$ such that

$$\inf_{x \in K} \langle \hat{\xi}, F_\xi(\bar{x})(x - G(\bar{x})) \rangle \geq \sup_{y \in -\text{int}(\mathbb{R}_+^s)} \langle \hat{\xi}, y \rangle. \quad (4)$$

If there exists $j \in [m]$ such that $\hat{\xi}_j < 0$ then $\sup_{y \in -\text{int}(\mathbb{R}_+^s)} \langle \hat{\xi}, y \rangle = +\infty$; then, the left hand of (4) equals to $+\infty$. This follows $K = \emptyset$, contrary to the assumption. Hence, $\hat{\xi} \in \Delta$ and $\hat{\xi}^T F_\xi(\bar{x})(x - G(\bar{x})) \geq 0$ for every $x \in K$. Therefore we obtain $\bar{x} \in \text{Sol}(\text{GVI}_\xi(F, G, K))$. The lemma is proved. $\square$

The main result in this section is stated established as follows.

Theorem 3.2. *Consider the problem $\text{GPVVI}(F, G, K)$ with $\text{Ker}(G^\infty) = \{0\}$. Suppose that the pair (G, K) has the property $\mathcal{A}$. Then, the following statements hold:*

(i) *If there exists $\xi \in \Delta$ such that*

$$\text{Sol}(F_\xi^\infty + \rho G^\infty, G^\infty, K^\infty) = \{0\} \quad (5)$$

for every $\rho \geq 0$ then $\text{GVI}_\xi(F, G, K)$ has a solution; hence, $\text{Sol}^w(F, G, K)$ is nonempty. In addition, if

$$\text{Sol}^w(F^\infty, G^\infty, K^\infty) = \{0\} \quad (6)$$

then $\text{Sol}^w(F, G, K)$ is a nonempty compact set;

(ii) *If there exists $\xi \in ri\Delta$ satisfying (5) for every $\rho \geq 0$ then $\text{Sol}(F, G, K)$ is nonempty.*

Proof. (i) Suppose that there exists $\xi \in \Delta$ satisfying (5) for every $\rho \geq 0$ and that the problem $\text{GVI}_\xi(F, G, K)$ has no solution. According to Lemma 2.1, there exists an exceptional family of elements for the pair (F_ξ, G) with respect to $0 \in \mathbb{R}^s$, i.e., there exist $\{x^k\} \subset \mathbb{R}^s$ satisfying $\|x^k\| \rightarrow \infty$ as $k \rightarrow \infty$ and $\alpha^k > 0$ such that $G(x^k) \in K$ and

$$\langle F_\xi(x^k) + \alpha^k G(x^k), y - G(x^k) \rangle \geq 0 \quad \forall y \in K.$$

It follows that

$$\left\langle \sum_{i=1}^m \xi_i F_i(x^k), y - G(x^k) \right\rangle + \alpha^k \langle G(x^k), y - G(x^k) \rangle \geq 0 \quad \forall y \in K. \quad (7)$$

By the fact that $\|x^k\| \rightarrow \infty$ as $k \rightarrow \infty$, we may assume that $\|x^k\| > 0$ for all $k \rightarrow \infty$ and $\frac{x^k}{\|x^k\|} \rightarrow \bar{h}$ for some $\bar{h} \in \mathbb{R}^s$ with $\|\bar{h}\| = 1$. Applying [24, Theorem 8.2] to $G(x^k) \in K$ and $\frac{1}{\|x^k\|^\sigma} \rightarrow 0$, we have

$$\frac{1}{\|x^k\|^\sigma} G(x^k) \rightarrow G^\infty(\bar{h}) \in K^\infty.$$

From the assumption that $\text{Ker}(G^\infty) = \{0\}$ it follows that $G^\infty(\bar{h}) \neq 0$. For some $y \in K$, dividing both sides of the inequality (7) by $\|x^k\|^{\sigma+\delta}$, we obtain that

$$A(x^k) + \alpha^k \|x^k\|^{\sigma-\delta} B(x^k) \geq 0, \quad (8)$$

where
$$A(x^k) := \left\langle \sum_{i=1}^m \frac{\xi_i F_i(x^k)}{\|x^k\|^\delta}, \frac{y - G(x^k)}{\|x^k\|^\sigma} \right\rangle \quad (9)$$

and
$$B(x^k) := \left\langle \frac{G(x^k)}{\|x^k\|^\sigma}, \frac{y - G(x^k)}{\|x^k\|^\sigma} \right\rangle \quad (10)$$

We have
$$\lim_{k \rightarrow \infty} B(x^k) = -\|G^\infty(\bar{h})\| < 0 \quad (11)$$

and
$$\lim_{k \rightarrow \infty} A(x^k) = \left\langle \sum_{i=1}^m \frac{\xi_i F_i(x^k)}{\|x^k\|^\delta}, \frac{y - G(x^k)}{\|x^k\|^\sigma} \right\rangle = -\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{h}), G^\infty(\bar{h}) \right\rangle. \quad (12)$$

We now show that $\{\rho^k := \alpha^k \|x^k\|^{\sigma-\delta}\}$ is bounded. Indeed, suppose, on the contrary, that $\rho^k \rightarrow +\infty$ as $k \rightarrow +\infty$. Then, dividing both sides of the equality (8) by ρ^k and letting $k \rightarrow \infty$ yields

$$-\|G^\infty(\bar{h})\| \geq 0,$$

contrary to (11). Hence $\{\rho^k\}$ is bounded. Without loss of generality, we may assume that $\rho^k \rightarrow \bar{\rho}$ for some $\bar{\rho} \in \mathbb{R}_+$. Fix $w \in K$. For every $h \in K^\infty$, we have $z := w + h\|x^k\|^\sigma \in K$. From (7) it follows that

$$\left\langle \sum_{i=1}^m \xi_i F_i(x^k) + \rho^k \|x^k\|^{-\sigma+\delta} G(x^k), w + h\|x^k\|^\sigma - G(x^k) \right\rangle \geq 0.$$

Dividing both sides of the latter inequality by $\|x^k\|^{\delta+\sigma}$ and letting $k \rightarrow +\infty$ yields:

$$\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{h}) + \bar{\rho} G^\infty(\bar{h}), h - G^\infty(\bar{h}) \right\rangle \geq 0.$$

This leads to $0 \neq \bar{h} \in \text{Sol}^w(F_\xi^\infty + \bar{\rho}G^\infty, G^\infty, K^\infty) = \{0\}$, a contradiction. Hence, $\text{GVI}_\xi(F, G, K)$ has a solution. From Lemma 3.1 it follows that $\text{Sol}^w(F, G, K)$ is nonempty.

Let any a sequence $\{z^k\} \subset \text{Sol}^w(F, G, K)$ such that $z^k \rightarrow \bar{z}$ as $k \rightarrow +\infty$ for some $\bar{z} \in \mathbb{R}^s$. For each k , there exists $\xi^k \in \Delta$ such that $z^k \in \text{Sol}^w(F_{\xi^k}, G, K)$; that is, for some $y \in K$,

$$G(z^k) \in K \quad \text{and} \quad \left\langle \sum_{i=1}^m \xi_i^k F_i(z^k), y - G(z^k) \right\rangle \geq 0. \quad (13)$$

From the boundedness of the sequence $\{\xi^k\}$ it follows that a subsequence $\{k_j\} \subset \{k\}$ such that $\xi^{k_j} \rightarrow \hat{\xi}$ as $j \rightarrow \infty$ for some $\hat{\xi} \in \Delta$. Passing the inequality (13) to limits as $k \rightarrow \infty$, we have

$$G(\bar{z}) \in K \quad \text{and} \quad \left\langle \sum_{i=1}^m \hat{\xi}_i F_i(\bar{z}), y - G(\bar{z}) \right\rangle \geq 0.$$

It follows that $\bar{z} \in \text{Sol}^w(F, G, K)$; hence, $\text{Sol}^w(F, G, K)$ is closed.

Finally, we show the boundedness of $\text{Sol}^w(F, G, K)$. Suppose, on the contrary, that $\text{Sol}^w(F, G, K)$ is unbounded. Then, there exists a sequence $\{y^k\} \subset \text{Sol}^w(F, G, K)$ such that $\|y^k\| \rightarrow +\infty$ as $k \rightarrow +\infty$. Without loss of generality we may assume that $\|y^k\| \rightarrow \infty$ as $k \rightarrow \infty$, $\|y^k\| \neq 0$ for all k , and $\|y^k\|^{-1}y^k \rightarrow \bar{v}$ with $\|\bar{v}\| = 1$. For each k , there exists $\xi^k \in \Delta$ such that $y^k \in \text{Sol}^w(F_{\xi^k}, G, K)$; that is, $G(y^k) \in K$ and

$$\left\langle \sum_{i=1}^m \xi_i^k F_i(y^k), y - G(y^k) \right\rangle \geq 0 \quad \forall y \in K. \quad (14)$$

Applying [24, Theorem 8.2] to $G(y^k) \in K$ and $\frac{1}{\|y^k\|^\sigma} \rightarrow 0$, we have

$$\frac{1}{\|y^k\|^{l-1}} G(y^k) \rightarrow G^\infty(\bar{v}) \in K^\infty$$

as $k \rightarrow \infty$. From the boundedness of the sequence $\{\xi^k\}$ it follows that a subsequence $\{k_j\} \subset \{k\}$ such that $\xi^{k_j} \rightarrow \bar{\xi}$ as $j \rightarrow \infty$ for some $\bar{\xi} \in \Delta$. Fix $w \in K$. For every $v \in K^\infty$, we have

$$z := w + v\|y^{k_j}\|^\sigma \in K.$$

From (14) it follows that

$$\left\langle \sum_{i=1}^m \xi_i^{k_j} F_i(y^{k_j}), w + v\|y^{k_j}\|^\sigma - G(y^{k_j}) \right\rangle \geq 0.$$

Dividing both sides of the latter inequality by $\|y^k\|^{\delta+\sigma}$ and letting $k \rightarrow \infty$ yields

$$\left\langle \sum_{i=1}^m \bar{\xi}_i F_i^\infty(\bar{v}), v - G^\infty(\bar{v}) \right\rangle \geq 0.$$

Then, $0 \neq \bar{v} \in \text{Sol}(F_\xi^\infty, G^\infty, K^\infty) \subset \text{Sol}^w(F^\infty, G^\infty, K^\infty)$, contrary to the assumption that $\text{Sol}^w(F^\infty, G^\infty, K^\infty) = \{0\}$. Therefore, $\text{Sol}^w(F, G, K)$ is bounded. Therefore, $\text{Sol}^w(F, G, K)$ is a compact set.

(ii) By similar arguments as in part (i), the nonemptiness of $Sol(F, G, K)$ is obvious. The proof is complete. $\square$

4. Some stability results

In this section, we study the upper/lower semicontinuity of the set-valued maps $Sol^w((\cdot, K))$ of GPVVI^w and $Sol((\cdot, K))$ of GPVVI with respect to the change in the problem parameters. Namely, we study the upper/lower semicontinuity of the set-valued maps

$$\Omega \ni \omega = (F', G') \mapsto Sol^w((F', G', K)),$$

$$\text{and} \quad \Omega \ni \omega = (F', G') \mapsto Sol((F', G', K)),$$

$$\text{where} \quad \Omega := \underbrace{\mathcal{P}^{[\delta, s]} \times \dots \times \mathcal{P}^{[\delta, s]}}_{m \text{ times}} \times \mathcal{P}^{[\sigma, s]}.$$

The following lemma is useful for proving the main theorem.

Lemma 4.1. [2, p.114] *Let I be an index set and let X and Y be topological spaces. Then, the union $\Phi = \cup_{i \in I} \Phi_i$ of a family of lower semicontinuous set-valued mappings Φ_i from X into Y is also a lower semicontinuous set-valued mapping from X into Y .*

A sufficient condition for the continuity of $Sol^w(\text{GPVVI}(\cdot, K))$ on Ω is proposed below.

Theorem 4.2. *Consider the problems $\text{GPVVI}(\omega, K)$ depending on the parameter $\omega = (F, G) \in \Omega$; $F = (F_1, \dots, F_m)$. Suppose that*

- (a) *the pair (G, K) has the property $\mathcal{A}$ and $\text{Ker}(G^\infty) = \{0\}$ for every $\omega \in \Omega$,*
- (b) *the condition (5) is satisfied for some $\xi \in \Delta$ and*
- (c) *F_i is strictly monotone respect to G on K for every $i \in [m]$ for every $\omega \in \Omega$.*

Then, the following statements hold:

- (i) *$Sol(\text{GPVI}_\xi(\cdot, K))$ is single-valued on Ω ;*
- (ii) *$Sol(\text{GPVI}_\xi(\cdot, K))$ is continuous on Ω ;*
- (iii) *$Sol^w(\text{GPVVI}(\cdot, K))$ is continuous on Ω .*

Proof. (i) By Theorem 3.2, for some $\bar{\omega} \in \Omega$, we obtain that $Sol(\text{GPVI}_\xi(\bar{\omega}, K))$ is nonempty. On the contrary, suppose that there exist $\bar{z}$ and $\bar{x}$ such that $\bar{z} \neq \bar{x}$ and $\{\bar{z}, \bar{x}\} \subset Sol(\text{GPVI}_\xi(\bar{\omega}, K))$. Then, we have $G(\bar{z}) \in K$, $G(\bar{x}) \in K$, and

$$\langle F_\xi(\bar{z}), G(\bar{x}) - G(\bar{z}) \rangle \geq 0 \quad \text{and} \quad \langle F_\xi(\bar{x}), G(\bar{z}) - G(\bar{x}) \rangle \geq 0.$$

It implies that $\langle F_\xi(\bar{x}) - F_\xi(\bar{z}), G(\bar{x}) - G(\bar{z}) \rangle \leq 0$. This contradicts the assumption that F_i is strictly monotone with respect to G on K for every $i \in [m]$. Therefore, $Sol(\text{GPVI}_\xi(\cdot, K))$ is single-valued on Ω .

(ii) We next show that the solution map $Sol(\text{GPVI}_\xi(\cdot, K))$ is lower semicontinuous on Ω . Indeed, on the contrary, suppose that there exists $\bar{\omega} \in \Omega$ such that $Sol(\text{GPVI}_\xi(\cdot, K))$ is not lower semicontinuous at $\bar{\omega}$. That is, there exists some

$\bar{x} \in \text{Sol}(\text{GPVI}_\xi(\bar{\omega}, K))$ and a sequence $\{\omega^l\} \subset \Omega$ satisfying $\omega^l \rightarrow \bar{\omega}$ such that, for any $z^l \in \text{Sol}(\text{GPVI}_\xi(\omega^l, K))$ satisfying $z^l \rightarrow \bar{z}$, one has $\bar{z} \neq \bar{x}$. By part (i), we have $\{\bar{x}\} = \text{Sol}(\text{GPVI}_\xi(\bar{\omega}, K))$. Since $z^l \in \text{Sol}(\text{GPVI}_\xi(\omega^l, K))$, we obtain that

$$G(z^l) \in K \text{ and } \langle F_\xi(z^l), z - G(z^l) \rangle \geq 0 \quad \forall z \in K. \quad (15)$$

For each $z \in K$, passing (15) to limits as $s \rightarrow \infty$ gives

$$G(\bar{z}) \in K \text{ and } \langle F_\xi(\bar{z}), z - G(\bar{z}) \rangle \geq 0.$$

This implies that $\bar{z} \in \text{Sol}(\text{GPVI}_\xi(\bar{\omega}, K))$. Hence, $\bar{z} = \bar{x}$, a contradiction. It follows that $\text{Sol}(\text{GPVI}_\xi(\cdot, K))$ is lower semicontinuous on Ω . Since the map $\text{Sol}(\text{GPVI}_\xi(\cdot, K))$ is single-valued on Ω , we conclude that $\text{Sol}(\text{GPVI}_\xi(\cdot, K))$ is continuous on Ω .

(iii) By Lemma 3.1, we have

$$\text{Sol}^w(\text{GPVVI}(\omega, K)) = \bigcup_{\xi \in \Delta} \text{Sol}(\text{GPVI}_\xi(\omega, K)).$$

Therefore, we deduce that $\text{Sol}^w(\text{GPVVI}(\cdot, K))$ is lower semicontinuous on Ω by using Lemma 4.1.

Finally, we show that $\text{Sol}^w(\text{GPVVI}(\cdot, K))$ is upper semicontinuous on Ω . Indeed, suppose that $\text{Sol}^w(\text{GPVVI}(\cdot, K))$ is not upper semicontinuous at $\bar{\omega}$ for some $\bar{\omega} \in \Omega$. Then, there exist an open U containing $\text{Sol}^w(\text{GPVVI}(\bar{\omega}, K))$, a sequence $\omega^l \rightarrow \bar{\omega}$ and $x^l \in \text{Sol}^w(\text{GPVVI}(\omega^l, K))$ such that $x^l \notin U$ for every l .

By the fact that $x^l \in \text{Sol}^w(\text{GPVVI}(\omega^l, K))$ and Lemma 3.1, there exist $\bar{\xi} \in \Delta$ such that $x^l \in \text{Sol}(\text{GPVI}_{\bar{\xi}}(\omega^l, K))$. From the assumption that F_i is strictly monotone respect to G on K for every $i \in [m]$ and the above part (i) it follows that $\text{Sol}(\text{GPVI}_{\bar{\xi}}(\cdot, K))$ is single-valued. Suppose that $\text{Sol}(\text{GPVI}_{\bar{\xi}}(\bar{\omega}, K)) = \{\bar{x}\}$. Then, $\text{Sol}(\text{GPVI}_{\bar{\xi}}(\cdot, K))$ is continuous at $\bar{\xi}$. Hence, $x^l \rightarrow \bar{x} \in \text{Sol}(\text{GPVI}_{\bar{\xi}}(\bar{\omega}, K)) \subset U$.

This contradicts the openness of the set U . Thus $\text{Sol}^w(\text{GPVVI}(\cdot, K))$ is upper semicontinuous on Ω and (iii) is shown. The proof is complete. $\square$

The following lemma is a Hartman-Stampacchia type theorem for the GPVVI.

Lemma 4.3. *Consider the problem $\text{GPVVI}(F, G, K)$ with $\text{Ker}(G^\infty) = \{0\}$. Suppose that G and K satisfy the property $\mathcal{A}$. Then, for each $\xi \in \Delta$, $\text{GPVI}_\xi(F, G, K)$ has a solution if K is a nonempty compact convex set.*

Proof. On the contrary, suppose that the problem $\text{GVI}_\xi(F, G, K)$ has no solution for some $\xi \in \Delta$. According to Lemma 2.1, there exists an exceptional family of elements for the pair (F_ξ, G) with respect to $0 \in \mathbb{R}^s$, i.e., there exist $\{x^k\} \subset \mathbb{R}^s$ satisfying $\|x^k\| \rightarrow \infty$ as $k \rightarrow \infty$ and $\alpha^k > 0$ such that $G(x^k) \in K$ and

$$-F_\xi(x^k) - \alpha^k G(x^k) \in \mathcal{N}_K(G(x^k)).$$

By the fact that $\|x^k\| \rightarrow \infty$ as $k \rightarrow \infty$, we may assume that $\|x^k\| > 0$ for all $k \rightarrow \infty$. Without loss of generality, we may assume that $\frac{x^k}{\|x^k\|} \rightarrow \bar{h}$ for some $\bar{h} \in \mathbb{R}^s$ with $\|\bar{h}\| = 1$. Applying [24, Theorem 8.2] to $G(x^k) \in K$ and $\frac{1}{\|x^k\|^\sigma} \rightarrow 0$, we have

$$\frac{1}{\|x^k\|^\sigma} G(x^k) \rightarrow G^\infty(\bar{h}) \in K^\infty = \{0\}.$$

From the assumption that $\text{Ker}(G^\infty) = \{0\}$ it follows that $\bar{h} = 0$, a contradiction. Therefore, $\text{GPVI}_\xi(F, G, K)$ has a solution. $\square$

Remark 4.4. Applying Lemma 4.3 to the case where $G = id_{\mathbb{R}^s}$, we get well-known Hartman-Stampacchia's theorem for the PVI (see [10]).

We obtain the following stability result.

Theorem 4.5. Suppose that G and K satisfy the property $\mathcal{A}$, and $\text{Ker}(G^\infty) = \{0\}$. The following statements are valid:

- (i) If F_i^∞ is strictly copositive with respect to G^∞ on K^∞ for every $i \in [m]$ then $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is a nonempty compact set for every $(\tilde{F}, \tilde{G}) \in \Omega$;
- (ii) If F_i^∞ is copositive with respect to G^∞ on K^∞ for every $i \in [m]$ and if there exists $\xi \in \Delta$ such that

$$\text{Sol}(F_\xi^\infty, G^\infty, K^\infty) = \{0\} \quad (16)$$

then $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is nonempty for every $(\tilde{F}, \tilde{G}) \in \Omega$. In addition, if (16) holds for every $\xi \in \Delta$ then $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is a compact set;

- (iii) If F_i is copositive with respect to G on K for every $i \in [m]$ then $\text{Sol}(F + c, G, K)$ is nonempty for every $c \in \mathbb{R}^{m \times n}$ satisfying

$$\sum_{i=1}^m \xi_i c_i \in \text{int}\{G^\infty(\text{Sol}^w(F_\xi^\infty, G^\infty, K^\infty))\}^* \quad (17)$$

for some $\xi \in \Delta$.

In addition, if (17) holds for every $\xi \in \Delta$ then $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is a compact set.

Proof. (i) For some $z^0 \in K$, for each $s = 1, 2, \dots$, denote

$$K_s = \{z \in \mathbb{R}^n : z \in K, \|z - z^0\| \leq s\}.$$

Then, we may assume that K_s is a nonempty compact set. By using Lemma 4.3 we obtain for some $(\tilde{F}, \tilde{G}) \in \mathcal{P}^{[m-1, n]} \times \mathcal{P}^{[l-1, n]}$ that $\text{GPVI}(F_\xi + \tilde{F}_\xi, G, K_s)$ has a solution for every s . Take any $x^s \in \text{Sol}(F_\xi + \tilde{F}_\xi, G + \tilde{G}, K_s)$. We prove that $\{x^s\}$ is bounded. Indeed, suppose, on the contrary, that $\{x^s\}$ is unbounded. Then, we may assume that $x^s > 0$ for every s and $x^s / \|x^s\| \rightarrow \bar{v}$ for some $\bar{v} \in \mathbb{R}^n$. By the fact that $x^s \in \text{Sol}(F_\xi + \tilde{F}_\xi, G + \tilde{G}, K_s)$, we have $G(x^s) + \tilde{G}(x^s) \in K_s$ and

$$\left\langle \sum_{i=1}^m \xi_i (F_i(x^s) + \tilde{F}_i(x^s)), z - G(x^s) - \tilde{G}(x^s) \right\rangle \geq 0 \quad (18)$$

for every $z \in K_s$. Since $G(x^s) + \tilde{G}(x^s) \in K$ and $\frac{1}{\|x^s\|^\sigma} \rightarrow 0$, applying [24, Theorem 8.2], we have

$$\frac{1}{\|x^s\|^\sigma} \left(G(x^s) + \tilde{G}(x^s) \right) \rightarrow G^\infty(\bar{v}) \in K^\infty$$

as $s \rightarrow \infty$.

Multiplying both sides of the inequality (18) by $\|x^s\|^{-(\delta+\sigma)}$ and taking $s \rightarrow \infty$ yields

$$\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), G^\infty(\bar{v}) \right\rangle \leq 0. \quad (19)$$

This contradicts the assumption that F_i^∞ is strictly copositive with respect to G^∞ on K^∞ for every $i \in [m]$. Therefore, $\{x^s\}$ is bounded. We may assume, without loss of generality, that $x^s \rightarrow \bar{x}$ for some $\bar{x} \in \mathbb{R}^n$. Since $G(x^s) + \tilde{G}(x^s) \in K$ and since K is closed, we have $G(\bar{x}) + \tilde{G}(\bar{x}) \in K$. Passing (18) to limits as $s \rightarrow \infty$, we obtain

$$\left\langle \sum_{i=1}^m \xi_i (F_i(\bar{x}) + \tilde{F}_i(\bar{x})), z - G(\bar{x}) - \tilde{G}(\bar{x}) \right\rangle \geq 0. \quad (20)$$

Hence, $\bar{x} \in \text{Sol}(F_\xi + \tilde{F}_\xi, G + \tilde{G}, K) \subset \text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$.

Suppose that $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is unbounded.

Then, there exists $\{y^s\} \subset \text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ such that we have $\|y^s\| \rightarrow \infty$ and $y^s/\|y^s\| \rightarrow \bar{y}$ for some $\bar{y} \in \mathbb{R}^n$. By Lemma 2.1, there exists $\xi^s \in \Delta$ such that $y^s \in \text{Sol}(F_{\xi^s} + \tilde{F}_{\xi^s}, G + \tilde{G}, K)$ for each s . By the fact that $\|\xi^s\| = 1$ for every s , we may assume that $\xi^s \rightarrow \hat{\xi}$ for some $\hat{\xi} \in \Delta$.

Repeating the above arguments, we obtain that $\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{y}), G^\infty(\bar{y}) \right\rangle \leq 0$. This contradicts the assumption that F_i^∞ is strictly copositive with respect to G^∞ on K^∞ for every $i \in [m]$. Thus, $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is bounded. The closedness of $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ follows from a similar argument as in the proof of Theorem 3.2 (part (i)). Therefore, $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is compact set.

(ii) By similar arguments as in part (i), we obtain (18) and (19). Since F_i^∞ is copositive with respect to G^∞ on K^∞ , we have

$$\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), G^\infty(\bar{v}) \right\rangle \geq 0.$$

By this and (19), we have

$$\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), G^\infty(\bar{v}) \right\rangle = 0. \quad (21)$$

For any $v \in K^\infty \setminus \{0\}$, one has $y^s := z^0 + \frac{\|G(x^s) + \tilde{G}(x^s)\|}{\|v\|} v \in K$ and

$$\|y^s - z^0\| = \|G(x^s) + \tilde{G}(x^s)\| \leq s.$$

Hence, $y^s \in K_s$. By (18), one has

$$\left\langle \sum_{i=1}^m \xi_i (F_i(x^s) + \tilde{F}_i(x^s)), \frac{\|G(x^s) + \tilde{G}(x^s)\|}{\|v\|} v - G(x^s) - \tilde{G}(x^s) \right\rangle \geq 0.$$

Multiplying both sides of the last inequality by $\|x^s\|^{-(\delta+\sigma)}$ and letting $s \rightarrow \infty$ yields

$$\left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), \frac{\|G^\infty(\bar{v})\|}{\|\bar{v}\|} \bar{v} \right\rangle \geq \left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), G^\infty(\bar{v}) \right\rangle = 0.$$

Hence, $\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), \bar{v} \rangle \geq 0$, that is, $\sum_{i=1}^m \xi_i F_i^\infty(\bar{v}) \in (K^\infty)^*$. By this and (21), we deduce that

$$G^\infty(\bar{v}) \in K^\infty, \quad \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}) \in (K^\infty)^* \quad \text{and} \quad \left\langle \sum_{i=1}^m \xi_i F_i^\infty(\bar{v}), G^\infty(\bar{v}) \right\rangle = 0,$$

that is, $0 \neq \bar{v} \in \text{Sol}(F_\xi^\infty, G^\infty, K^\infty)$, (22)

contrary to the assumption that $\text{Sol}(F_\xi^\infty, G^\infty, K^\infty) = \{0\}$. Therefore, $\{x^s\}$ is bounded. The nonemptiness of $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ follows from a similar argument as in part (i).

Repeating the above arguments as in part (i), we obtain that

$$0 \neq \bar{y} \in \text{Sol}(F_\xi + \tilde{F}_\xi, G + \tilde{G}, K),$$

contrary to the assumption that (16) holds for every $\xi \in \Delta$.

Thus, $\text{Sol}^w(F + \tilde{F}, G + \tilde{G}, K)$ is bounded.

(iii) Let $c \in \mathbb{R}^{m \times n}$ satisfying $\sum_{i=1}^m \xi_i c_i \in \text{int}\{G^\infty(\text{Sol}(F_\xi^\infty, G^\infty, K^\infty))\}^*$. By similar arguments as in part (i) with $\tilde{F} \equiv c \in \mathbb{R}^m$ and $\tilde{G} \equiv 0$, we obtain (18)–(21). Repeating the arguments of part (i), one gets $\bar{v} \in \text{Sol}(F_\xi^\infty, G^\infty, K^\infty)$. From (18), taking $z = 0$, we have

$$\left\langle \sum_{i=1}^m \xi_i (F_i(x^s) + c_i), -G(x^s) \right\rangle \geq 0.$$

This implies that

$$\left\langle \sum_{i=1}^m \xi_i c_i, G(x^s) \right\rangle \leq - \left\langle \sum_{i=1}^m \xi_i F_i(x^s), G(x^s) \right\rangle \leq 0,$$

since F_i is copositive with respect to G on K for every $i \in [m]$. Dividing both sides of the last inequality by $\|x^s\|^\sigma$ and letting $s \rightarrow \infty$ yields

$$\left\langle \sum_{i=1}^m \xi_i c_i, G^\infty(\bar{v}) \right\rangle \leq 0,$$

contrary to the assumption that $\sum_{i=1}^m \xi_i c_i \in \text{int}\{G^\infty(\text{Sol}(F_\xi^\infty, G^\infty, K^\infty))\}^*$.

Therefore, $\text{Sol}^w(F + c, G, K)$ is nonempty.

The boundedness and closedness of $\text{Sol}^w(F + c, G, K)$ follows from a similar analysis as in part (i). □

By Theorem 4.5, we get the following corollary.

Corollary 4.6. *Suppose that the assumptions in Theorem 4.5 are satisfied. The following statements are valid:*

- (i') *If F_i^∞ is strictly copositive with respect to G^∞ on K^∞ for every $i \in [m]$ then $\text{Sol}(F + \tilde{F}, G + \tilde{G}, K)$ is nonempty for every $(\tilde{F}, \tilde{G}) \in \mathcal{P}^{[m-1, n]} \times \mathcal{P}^{[l-1, n]}$;*
- (ii') *If F_i^∞ is copositive with respect to G^∞ on K^∞ for every $i \in [m]$ and (16) holds for some $\xi \in \text{ri}\Delta$ then $\text{Sol}(F + \tilde{F}, G + \tilde{G}, K)$ is nonempty for every $(\tilde{F}, \tilde{G}) \in \mathcal{P}^{[m-1, n]} \times \mathcal{P}^{[l-1, n]}$.*
- (iii') *If F_i is copositive with respect to G on K for every $i \in [m]$ then, for every $c \in \mathbb{R}^{m \times n}$ satisfying (17) for some $\xi \in \text{ri}\Delta$, the set $\text{Sol}(F + c, G, K)$ is nonempty.*

Proof. The desired conclusion follows from a similar argument as in the proof of Theorem 4.5. $\square$

5. Applications for polynomial variational inequality

Recall that if $G = \text{id}_{\mathbb{R}^s}$, $m = 1$, and $F = F_1$ then $\text{GPVVI}(F, G, K)$ reduces to the following *polynomial variational inequality* (PVI) (see [5, 10])

$$\text{Find } \bar{x} \in K \text{ s.t. } \langle F(\bar{x}), x - \bar{x} \rangle \geq 0 \quad \forall x \in K. \quad (\text{PVI}(F, K))$$

Applying Theorem 4.5(ii) to the problem $(\text{PVI}(F, K))$, we obtain the following corollary.

Corollary 5.1. *Consider the problem $(\text{PVI}(F, K))$ with K being nonempty. Then, the following statements are valid:*

- (a) *If F^∞ is strictly copositive on K^∞ then $\text{Sol}(F + \tilde{F}, K)$ is a nonempty compact set for every $\tilde{F} \in \mathcal{P}^{[m-1, n]}$;*
- (b) *If F^∞ is copositive on K^∞ and $\text{Sol}(F^\infty, K^\infty) = \{0\}$ then $\text{Sol}(F + \tilde{F}, K)$ is a nonempty compact set for every $\tilde{F} \in \mathcal{P}^{[m-1, n]}$;*
- (c) *If F^∞ is copositive on K^∞ then $\text{Sol}(F + c, K)$ is a nonempty compact set for every $c \in \mathbb{R}^{m \times n}$ satisfying $c \in \text{int}\{\text{Sol}(F^\infty, K^\infty)\}^*$.*

Remark 5.2. Corollary 5.1(c) is an extension of [8, Theorem 4.1] because the assumption “ $0 \in K$ ” is omitted. The obtained results in Corollary 5.1 are also generalizations of Theorems 6.3–6.5 in [11]. Corollary 5.1 gives a partial answer to Question 5.2 in [28]. Theorem 6.2 in [6] gives a result on the solution existence for weakly homogeneous variational inequality under the assumption $\text{int}(K^*) \neq \emptyset$. However, this condition is omitted in Corollary 5.1.

6. Conclusions

In the presented paper, sufficient conditions for the existence of Pareto solutions of the GPVVI are established (Theorem 3.2). Due to this result, we present a sufficient condition for upper/lower continuity of the weak Pareto solution map of the GVVI (Theorem 4.2). Some stability results for the GPVVI are also proposed (Theorem 4.5). Our results develop and complement the previous ones for VIs, GPVIs and VVIs in [8, 11, 15, 26, 28, 31].

Acknowledgments. A part of this work was done at Vietnam Institute for Advanced Study in Mathematics (VIASM). The authors would like to thank the VIASM for hospitality during our recent stay at the Institute. This research is funded by Hanoi Pedagogical University 2 under Grant Number HPU2.2023-UT-06.

References

- [1] X. Bai, M. Zheng, Z. H. Huang: *Unique solvability of weakly homogeneous generalized variational inequalities*, J. Glob. Optim. 80/4 (2021) 921–943.
- [2] C. Berge: *Topological Spaces*, Oliver and Boyd, London (1963).
- [3] F. Giannessi: *Theorems of alternative, quadratic programs and complementarity problems*, in: *Variational Inequality and Complementarity Problems*, R. W. Cottle et al. (eds.), Wiley, New York (1980) 151–186.
- [4] F. Giannessi (ed.): *Vector Variational Inequalities and Vector Equilibria*, Kluwer Academic Publishers, Dordrecht (2000).
- [5] M. S. Gowda: *Polynomial complementarity problems*, Pac. J. Optimization 13 (2017) 227–241.
- [6] M. S. Gowda, D. Sossa: *Weakly homogeneous variational inequalities and solvability of nonlinear equations over cones*, Math. Program. 177/1-2 (2019) 149–171.
- [7] J. Y. Han, Z. H. Huang, S. C. Fang: *Solvability of variational inequality problems*, J. Optim. Theory Appl. 122/3 (2004) 501–520.
- [8] V. T. Hieu: *Solution maps of polynomial variational inequalities*, J. Global Optim. 77 (2020) 807–824.
- [9] N. T. T. Huong, P. D. Khanh, N. D. Yen: *Multivalued Tikhonov trajectories of general affine variational inequalities*, J. Optim. Theory Appl. 158/1 (2013) 85–96.
- [10] D. Kinderlehrer, G. Stampacchia: *An Introduction to Variational Inequalities and Their Applications*, Academic Press, New York (1980).
- [11] G. M. Lee, N. N. Tam, N. D. Yen: *Quadratic Programming and Affine Variational Inequalities: A Qualitative Study*, Nonconvex Optimization and its Application Vol. 78, Springer, New York (2005).
- [12] L. Ling, C. Ling, H. He: *Properties of the solution set of generalized polynomial complementarity problems*, Pac. J. Optim. 16 (2020) 155–174.
- [13] X. P. Luo: *Tikhonov regularization methods for inverse variational inequalities*, Optim. Letters 8/3 (2014) 877–887.
- [14] X. P. Luo, J. Yang: *Regularization and iterative methods for monotone inverse variational inequalities*, Optim. Letters 8/4 (2014) 1261–1272.
- [15] T. V. Nghi: *On stability of solutions to parametric generalized affine variational inequalities*, Optimization 67/2 (2018) 269–285.
- [16] T. V. Nghi, N. N. Tam: *Continuity and directional differentiability of the optimal value function in parametric quadratically constrained nonconvex quadratic programs*, Acta Math. Vietnam. 42 (2017) 311–336.
- [17] T. V. Nghi, N. N. Tam: *Stability for parametric extended trust region subproblems*, Pac. J. Optim. 15 (2019) 111–129.
- [18] T. V. Nghi, N. N. Tam: *A Frank-Wolfe-type theorem for cubic programs and solvability for quadratic variational inequalities*, J. Optim. Theory Appl. 187 (2020) 448–468.

- [19] T. V. Nghi, N. N. Tam: *General variational inequalities: solution existence, Tikhonov-type regularization, and well-posedness*, Acta Math. Vietnam. 47 (2022) 539–552.
- [20] T. V. Nghi, N. N. Tam, V. T. Anh: *On generalized polynomial variational inequality. Part 1: Existence of solutions*, Pac. J. Optim. 17/1 (2021) 115–132.
- [21] M. A. Noor: *Quasi variational inequalities*, Appl. Math. Letters 1/4 (1988) 367–370.
- [22] M. A. Noor: *Extended general variational inequalities*, Appl. Math. Letters 22/2 (2009) 182–186.
- [23] J. S. Pang, J. C. Yao: *On a generalization of a normal map and equation*, SIAM J. Control Optim. 33/1 (1995) 168–184.
- [24] R. T. Rockafellar: *Convex Analysis*, Princeton University Press, Princeton (1970).
- [25] G. Stampacchia: *Formes bilinéaires coercitives sur les ensemble convexes*, C. R. Acad. Sci. Paris 258 (1964) 4413–4416.
- [26] N. N. Tam: *Some stability results for the semi-affine variational inequality problem*, Acta Math. Vietnam. 29 (2004) 271–280.
- [27] N. N. Tam, T. V. Nghi: *On the solution existence and stability of quadratically constrained nonconvex quadratic programs*, Optim. Lett. 12 (2018) 1045–1063.
- [28] Y. Wang, Z. H. Huang, L. Qi: *Global uniqueness and solvability of tensor variational inequalities*, J. Optim. Theory Appl. 177 (2018) 137–152.
- [29] J. Wang, Z. H. Huang, Y. Xu: *Existence and uniqueness of solutions of the generalized polynomial variational inequality*, Optim. Lett. 14 (2020) 1571–1582.
- [30] N. D. Yen: *An introduction to vector variational inequalities and some new results*, Acta Math. Vietnam. 41/3 (2016) 505–529.
- [31] N. D. Yen, J. C. Yao: *Monotone affine vector variational inequalities*, Optimization 60/1-2 (2011) 53–68.