

Variational Principles for Nonautonomous Dynamical Systems

Andrzej Biś^{1,*}

1. Faculty of Mathematics and Computer Science, University of Łódź, Poland

Abstract: We study thermodynamical formalism of a discrete nonautonomous dynamical system determined by a sequence of continuous self-maps of a compact metric space. Using the methods of Convex Analysis we get variational principles for pressure functions determined by the system.

Keywords: Finitely additive set function, pressure function, variational principle, nonautonomous dynamical systems.

2010 Mathematics Subject Classification: Primary: 37D35, 37C85; secondary: 43A05.

1. Introduction and structure of the paper

In 1959, Kolmogorov and Sinai developed the metric entropy from Shannon's information theory into ergodic theory. Since then, the notion of entropy has played a crucial role in the study of dynamical systems. In 1965, the notion of topological entropy, denoted here by $h_{\text{top}}(f)$, of a classical dynamical system, determined by a single continuous self-map $f : X \rightarrow X$ of a compact topological space X , was introduced by Adler, Konheim and McAndrew [1] as a topological conjugacy invariant. In 1971, Bowen [6] and Dinaburg [9] gave a new, but equivalent, definition in the case when the space under consideration is metrizable.

Another approach which resembles the construction of Hausdorff measure and Hausdorff dimension was used by Bowen [7] to extend the concept of topological entropy for noncompact sets. This approach was developed by Pesin and Pitskel [20] and presented in details by Pesin [19] who elaborated a dimension theory of Carathéodory structures.

Due to the Krylov-Bogoliubov Theorem [8] for a continuous map $f : X \rightarrow X$ there exists a Borel probability f -invariant measure μ , defined on a Borel σ -algebra on X . Therefore, the space of $\mathcal{P}_f(X)$ of f -invariant probability measures defined on the σ -algebra $\mathfrak{B}(X)$ of the Borel subsets of X is not empty. Thus, one can define and study measure-theoretic entropy, denoted here by $h_\mu(f)$, of a continuous transformation f with respect to a measure $\mu \in \mathcal{P}_f(X)$. The famous variational principle relates topological entropy of a continuous self-map $f : X \rightarrow X$, acting on a compact metric space X , to its measure-theoretic entropy.

It says that
$$h_{\text{top}}(f) = \sup\{h_\mu(f) : \mu \in \mathcal{P}_f(X)\}.$$

Equivalently, it can be rewritten for the topological pressure $P_{\text{top}}(f, \varphi)$ of the map f (Theorem 9.10 in [22]) as

$$P_{\text{top}}(f, \varphi) = \sup_{\mu \in \mathcal{P}_f(X)} \left\{ h_\mu(f) + \int_X \varphi d\mu \right\}, \quad (1)$$

where $\varphi : X \rightarrow \mathbb{R}$ is a continuous potential.

Also, it is known a dual variational principle (Theorem 9.12 in [22]) for a continuous self-map $f : X \rightarrow X$, with finite topological entropy. It expresses the measure-theoretic entropy $h_{\mu_0}(f)$ of f through the topological pressure $P_{\text{top}}(f, \varphi)$, it says that $h_{\mu_0}(f) = \inf\{P_{\text{top}}(f, \varphi) - \int_X \varphi d\mu_0 : \varphi \in C(X)\}$ if and only if $h_{\mu_0}(f)$ is upper semi-continuous at μ_0 , where $C(X)$ is the space of all continuous maps $\varphi : X \rightarrow \mathbb{R}$.

*Corresponding author.

E-mail addresses: A. Biś*: andrzej.bis@wmii.uni.lodz.pl

<https://doi.org/10.68381/jca34017>

Received date: October 31, 2025; Accepted date: April 2, 2026

© 2027 Published by Banach Press LLC. (Dedicated to Alexander Plakhov on the occasion of his 65th birthday.)

One of the aims of the thermodynamic formalism is to establish the existence of measures with maximal entropy. Such measures, called equilibrium states, include as specific examples Gibbs measures and physical measures. With some additional assumptions a classical dynamical system admits a measure of maximal entropy.

There are several proposals to extend the notions of measure-theoretic entropy and topological pressure to more general settings. One of them are nonautonomous discrete dynamical systems (in short: NADDS), determined by sequences of continuous self-maps $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$, which are natural generalization of a classical dynamical system.

Kolyada and Snoha [15] introduced and studied the notion of topological entropy of NADDS. Many properties for such dynamical systems have been studied for many years (cf. [2], [11], [12], [14], [15], [18], [23]). For example, Huang, Wen and Zeng [11] introduced and studied the topological pressure of NADDS. Equivalent definitions of topological pressures and basic properties of these notions one can find in [16].

A main difficulty in setting up the variational principle for NADDS is a lack of common invariant measures; there is no counterpart to the Krylov-Bogoliubov Theorem.

Kawan [12] considers at the same time a sequence of probability spaces $X_{1,\infty} := \{X_n, \mu_n\}_{n \in \mathbb{N}}$ and a sequence of continuous maps $f_{1,\infty} := \{f_n : X_n \rightarrow X_{n+1}\}_{n \in \mathbb{N}}$ defined on the sequence of compact metric spaces $\{X_n\}_{n \in \mathbb{N}}$ with a sequence of measures $\mu_{1,\infty} := \{\mu_n - \text{measure on } X_n\}_{n \in \mathbb{N}}$. The sequence of measures $\mu_{1,\infty}$ is called $f_{1,\infty}$ -invariant if $(f_n^*)\mu_n = \mu_{n+1}$, for any $n \in \mathbb{N}$. Let $\mathcal{P}_n$ be a finite measurable partition of X_n and let $\mathcal{P}_{1,\infty} := \{\mathcal{P}_n\}_{n \in \mathbb{N}}$ be the sequence of all finite partitions. Recall that the entropy of a finite measurable partition $P = \{P_1, \dots, P_m\}$ of a probability space $(X, \mathcal{A}, \mu)$, where $\mathcal{A}$ is a σ -algebra of X , is defined as $H_\mu(P) := -\sum_{i=1}^m \mu(P_i) \log(\mu(P_i))$. Kawan defines topological entropy $h_{top}(f_{1,\infty})$ in the same way as Kolyada and Snoha in [15]. Since the measure entropy of a partition $\mathcal{P}_{1,\infty}$ defined as follows

$$h_{\mu_1}(f_{1,\infty}, \mathcal{P}_{1,\infty}) := \limsup_{n \rightarrow \infty} \frac{1}{n} H_{\mu_1} \left(\bigvee_{i=0}^{n-1} f_1^{-i} \mathcal{P}_{i+1} \right)$$

can be infinite even for simple systems (see Example 14 and Example 18 in [12]) he chooses a sufficiently nice subclass $\mathcal{E}_{max}$ from the class of all sequences $\mathcal{P}_{1,\infty}$, called maximal admissible class (for definition see Subsection 3.2 of [12]), to get a reasonable notion of metric entropy. But to get partial variational principle he even chooses a subclass $\mathcal{E}_M \subset \mathcal{E}_{max}$, called Misiurewicz class, described in Subsection 4.1 of [12], to define measure entropy

$$h_{\mathcal{E}_M}(f_{1,\infty}) := \sup_{\mathcal{P}_{1,\infty} \in \mathcal{E}_M} h_{\mu_1}(f_{1,\infty}, \mathcal{P}_{1,\infty}).$$

His main result (Theorem 28 in [12]) is a partial variational principle. It ensures that for an equicontinuous topological nonautonomous dynamical system $(X_{1,\infty}, f_{1,\infty})$, with $f_{1,\infty}$ -invariant sequence $\mu_{1,\infty}$, the inequality $h_{\mathcal{E}_M}(f_{1,\infty}) \leq h_{top}(f_{1,\infty})$ holds.

Local measure entropies, in sense of Brin and Katok [8], are well defined for discrete nonautonomous dynamical systems. In this context, there were several approaches to relate the topological entropy of a system to its local measure entropies.

For example, some generalization of the result of Kawan [12] was given by Biś [2] who applied local lower measure entropy $h_\mu^{Low}(f_{1,\infty})(x)$ and local upper measure entropy $h_\mu^{Upp}(f_{1,\infty})(x)$ of the system at a point $x \in X$, with respect to a Borel probability measure μ on X .

Theorem 1 in [2] says that for a Borel probability measure μ on X a subset $E \subset X$ and $s > 0$ one has: If for any point $x \in E$ the inequality $h_\mu^{Low}(f_{1,\infty})(x) \geq s$ holds and $\mu(E) > 0$, then $h_{top}(f_{1,\infty}, E) \geq s$.

Feng and Huang [10] proved that the topological entropy of NADDS coincides with the supremum of integrals of lower local measure entropy function, where the supremum is taking over all Borel probability measures with full support.

In the paper, we prove that for the topological pressure of a nonautonomous dynamical system the abstract variational principle and dual abstract variational principle (Theorem A) hold. In Theorem B, assuming that the classical variational principle holds for topological pressure, we show the relations between different measure entropies and invariant measures of the system. Next, we use Misiurewicz construction to introduce Misiurewicz pressure and Misiurewicz entropy. In Theorem C we prove that for Misiurewicz pressure the abstract variational principle and abstract dual variational principle hold as well. In Theorem D, assuming that the classical variational principle holds for Misiurewicz pressure, we show the relations between different measure entropies and invariant measures of the system.

The paper is organized as follows. In Section 2, we introduce Convex Analysis approach to NSDDS. In particular we present necessary notions and ideas to formulate Theorem A on abstract variational principle for the topological pressure of a nonautonomous dynamical system $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$. We show that topological pressure determines a pressure function. The variational principle stated in Theorem A for topological pressure ensures that there always exist measures for which the right-hand side of (2) attains the maximum. In Corollary 2.1 we state that such a measure is unique. We define the space $\mathcal{P}_{f_{1,\infty}}(X)$ of $f_{1,\infty}$ -invariant measures and provide examples (Example 2.2 and Example 2.3) of nonautonomous dynamical systems for which this space $\mathcal{P}_{f_{1,\infty}}(X)$ is not empty. In Theorem B, under the assumption that for a nonautonomous dynamical system $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$ we have also classical variational principle for topological pressure, we present relations between different entropies of the system. In particular we prove that $f_{1,\infty}$ -invariant measures are determined by the positivity of the abstract measure entropy. Next we define Misiurewicz entropy and Misiurewicz pressure of a nonautonomous dynamical system $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$. We show that Misiurewicz pressure determines a pressure function. We state Theorem C, expressed for the Misiurewicz pressure, which is an analogue of Theorem A expressed for the topological pressure of a nonautonomous dynamical system. The variational principle stated in Theorem C for Misiurewicz pressure ensures that there always exist measures for which the right-hand side of (5) attains the maximum. In Corollary 2.4 we state that such a measure is unique. Next we state Theorem D, expressed for the Misiurewicz pressure, which is an analogue of Theorem B expressed for the topological pressure.

Section 3 is devoted to the main result of Biś et al. [3], where the authors used methods from Convex Analysis to obtain upper semi-continuous entropy-like maps adapted to any pressure function on a suitable Banach space, and without summoning any dynamics. An abstract pressure function (Definition 3.1) defined on a Banach space is introduced. In Theorem 3.2 stated for an abstract pressure function we obtain an abstract measure entropy, abstract variational principle and abstract dual variational principle.

In Section 4, we introduce in details the notion of topological pressure of NADDS, apply the results of [3], and provide proofs of Theorem A and Theorem B. In Section 5, we introduced Misiurewicz pressure of a nonautonomous dynamical system, based on [17], and present proofs of Theorem C and Theorem D.

2. Main results

In this self-contained paper we present two different definitions of pressures of NADDS determined by continuous self-maps $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$. The first one is called the topological pressure (see: Definition 4.1) and the second one is called the Misiurewicz pressure (see: Definition 5.1). To both pressures we apply the main results of [3], where the authors used methods from Convex Analysis to obtain upper semi-continuous entropy-like maps adapted to any abstract pressure function on a suitable Banach space, and without summoning any dynamics.

For a compact metric space (X, d) we denote by $P(X)$ the space of all Borel probability measures on X , and by $C(X)$ the space of all continuous maps $\varphi : X \rightarrow \mathbb{R}$, called potentials. Let $P_{\text{top}}(f_{1,\infty}, \varphi)$ (resp., $h_{\text{top}}(f_{1,\infty})$) denote the topological pressure (resp., the topological entropy) a NADDS $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$, with respect to the potential $\varphi \in C(X)$. Applying the strategy of [3] to topological pressure of NADDS we get the following abstract variational principle and abstract dual variational principle.

Theorem A. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system defined on a compact metric space (X, d) with $h_{\text{top}}(f_{1,\infty}) < +\infty$. Then there exists an upper semi-continuous function $\mathfrak{h}_{f_{1,\infty}} : \mathcal{P}(X) \rightarrow \mathbb{R}$ such that

$$P_{\text{top}}(f_{1,\infty}, \varphi) = \max_{\mu \in \mathcal{P}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int_X \varphi d\mu \right\} \quad \forall \varphi \in C(X) \quad (2)$$

and

$$\mathfrak{h}_{f_{1,\infty}}(\mu) = \inf_{\varphi \in C(X)} \left\{ P_{\text{top}}(f_{1,\infty}, \varphi) - \int_X \varphi d\mu \right\} \quad \forall \mu \in \mathcal{P}(X). \quad (3)$$

Corollary 2.1. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system with finite topological entropy $h_{\text{top}}(f_{1,\infty})$, and a continuous potential $\varphi \in C(X)$. Then there exists a unique measure $\mu \in P(X)$ such that $P_{\text{top}}(f_{1,\infty}, \varphi) = h_{f_{1,\infty}}(\mu) + \int_X \varphi d\mu$.

In the next theorem we introduce a new entropy like quantity $h_{f_{1,\infty}}^*(\mu)$ (for definition see Subsection 4.4) for a NADDS and we present some properties of an abstract measure-entropy of nonautonomous dynamical systems.

Let $\mathcal{P}_{f_{1,\infty}}(X)$ be the space of $f_{1,\infty}$ -invariant measures, i.e.

$$\mathcal{P}_{f_{1,\infty}}(X) := \{\mu \in \mathcal{P}(X) : \int (\psi \circ f_n) d\mu = \int \psi d\mu, \forall n \in \mathbb{N}, \psi \in C(X)\}.$$

Below we present examples of NADDS admitting common invariant measures.

Example 2.2. Assume that (X, d) is a compact metric space and there are two commuting continuous maps $f, g : X \rightarrow X$, that are different from the identity map on X . Then, a Borel probability measure μ which is f -invariant, is g -invariant as well (see [21] p. 98). So, for $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ consisting of pairwise commuting maps (i.e. $f_n \circ f_m = f_m \circ f_n$ for any $m, n \in \mathbb{N}$) there exists a common $f_{1,\infty}$ -invariant measure ν and $\mathcal{P}_{f_{1,\infty}}(X) \neq \emptyset$. $\square$

Example 2.3. The orthogonal group $O(m)$ acting on the m -dimensional sphere S^m is a non-abelian group admitting $O(m)$ -invariant Haar measure. Therefore, for any $f_{1,\infty} = \{f_n : S^m \rightarrow S^m\}_{n=1}^\infty$, with $f_n \in O(m)$ for any $n \in \mathbb{N}$, the Haar measure is $f_{1,\infty}$ -invariant and $\mathcal{P}_{f_{1,\infty}}(X) \neq \emptyset$. $\square$

Theorem B. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system defined on a compact metric space (X, d) . Let $P_{\text{top}}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ be the pressure function and $h_{f_{1,\infty}} : \mathcal{P}(X) \rightarrow \mathbb{R}$ its variational metric entropy given by the equality (3). Assume that $\mathcal{P}_{f_{1,\infty}}(X) \neq \emptyset$ and $P_{\text{top}}(f_{1,\infty}, \cdot)$ also satisfies a variational principle

$$P_{\text{top}}(f_{1,\infty}, \varphi) = \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} \quad \forall \varphi \in C(X) \quad (4)$$

with respect to a non-negative metric entropy map $h_{f_{1,\infty}} : \mathcal{P}_{f_{1,\infty}}(X) \rightarrow [0, +\infty[$.

Then

(a) If $\varphi \in C(X)$ and $\mu_\varphi \in \mathcal{P}(X)$ attains the maximum in (2), then μ_φ is $f_{1,\infty}$ -invariant, that is,

$$\int (\psi \circ f_n) d\mu_\varphi = \int \psi d\mu_\varphi \quad \forall n \in \mathbb{N} \quad \forall \psi \in C(X).$$

(b) Given $\mu \in \mathcal{P}(X)$, one has $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$ if and only if $h_{f_{1,\infty}}(\mu) \geq 0$.

(c) For every $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$, one has $0 \leq h_{f_{1,\infty}}(\mu) \leq h_{f_{1,\infty}}^*(\mu) \leq h_{f_{1,\infty}}(\mu)$.

(d) For every $\varphi \in C(X)$

$$P_{\text{top}}(f_{1,\infty}, \varphi) = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^*(\mu) + \int \varphi d\mu \right\}.$$

In the second part of the paper we consider Misiurewicz pressure $P_{\text{Mis}}(f_{1,\infty}, \varphi)$ (see Definition 5.1) and Misiurewicz entropy $h_{\text{Mis}}(f_{1,\infty})$ of $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$, defined by the equality $h_{\text{Mis}}(f_{1,\infty}) := P_{\text{Mis}}(f_{1,\infty}, \phi_0)$, where $\phi_0 \equiv 0$. We get the following results, similar to Theorem A and Theorem B.

Theorem C. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system defined on a compact metric space (X, d) , with $h_{\text{Mis}}(f_{1,\infty}, X) < +\infty$. Then there exists an upper semi-continuous function $h_{f_{1,\infty}}^{\text{Mis}} : \mathcal{P}(X) \rightarrow \mathbb{R}$ such that

$$P_{\text{Mis}}(f_{1,\infty}, \varphi) = \max_{\mu \in \mathcal{P}(X)} \left\{ h_{f_{1,\infty}}^{\text{Mis}}(\mu) + \int_X \varphi d\mu \right\} \quad \forall \varphi \in C(X) \quad (5)$$

and

$$\mathfrak{h}_{f_{1,\infty}}^{Mis}(\mu) = \inf_{\varphi \in C(X)} \left\{ P_{Mis}(f_{1,\infty}, \varphi) - \int \varphi d\mu \right\} \quad \forall \mu \in \mathcal{P}(X). \quad (6)$$

Corollary 2.4. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system with $h_{Mis}(f_{1,\infty}) < \infty$ and a continuous potential $\varphi \in C(X)$ there exists a unique measure $\mu_1 \in P(X)$ such that $P_{Mis}(f_{1,\infty}, \varphi) = \mathfrak{h}_{f_{1,\infty}}^{Mis}(\mu_1) + \int_X \varphi d\mu_1$.

In the next theorem we introduce a new entropy-like quantity $h_{f_{1,\infty}}^{*Mis}$ (see Subsection 5.5) for a NADDS and we present some properties of an abstract measure-entropy of $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$.

Theorem D. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system defined on a compact metric space (X, d) . Let $P_{Mis}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ be the Misiurewicz pressure function and $\mathfrak{h}_{f_{1,\infty}}^{Mis} : \mathcal{P}(X) \rightarrow \mathbb{R}$ denote its variational metric entropy given by the equality (6). Assume that $\mathcal{P}_{f_{1,\infty}}(X) \neq \emptyset$ and $P_{Mis}(f_{1,\infty}, \cdot)$ also satisfies a variational principle

$$P_{Mis}(f_{1,\infty}, \varphi) = \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^{Mis}(\mu) + \int \varphi d\mu \right\} \quad \forall \varphi \in C(X) \quad (7)$$

with respect to a non-negative metric entropy map $h_{f_{1,\infty}}^{Mis} : \mathcal{P}_{f_{1,\infty}}(X) \rightarrow [0, +\infty[$.

Then

(a) If $\varphi \in C(X)$ and $\mu_\varphi \in \mathcal{P}(X)$ attains the maximum in (5), then μ_φ is $f_{1,\infty}$ -invariant, that is,

$$\int (\psi \circ f_n) d\mu_\varphi = \int \psi d\mu_\varphi \quad \forall n \in \mathbb{N} \quad \forall \psi \in C(X).$$

(b) Given $\mu \in \mathcal{P}_a(X)$, one has $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$ if and only if $\mathfrak{h}_{f_{1,\infty}}^{Mis}(\mu) \geq 0$.

(c) For every $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$, one has $0 \leq h_{f_{1,\infty}}^{Mis}(\mu) \leq h_{f_{1,\infty}}^{*Mis}(\mu) \leq \mathfrak{h}_{f_{1,\infty}}^{Mis}(\mu)$.

(d) For every $\varphi \in C(X)$

$$P_{Mis}(f_{1,\infty}, \varphi) = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}^{Mis}(\mu) + \int \varphi d\mu \right\} = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^{*Mis}(\mu) + \int \varphi d\mu \right\}.$$

Remark 2.5. In context of group actions or semigroup actions results similar to Theorem B or Theorem D one can find in [4]. $\square$

3. Abstract variational principle

This section is on the main result of Biś et al. [3], where the authors used methods from Convex Analysis to obtain upper semi-continuous entropy-like maps adapted to any pressure function on a suitable Banach space, and without summoning any dynamics. Thus, this strategy is unspecific enough to be applied to either the classical topological pressure associated to a single dynamics (as done in [3]) or to the recently defined notions of topological pressure and Misiurewicz pressure for nonautonomous dynamical systems, we address in this work. Such an abstract variational principle can be applied to obtain variational principles using either the space of probability measures or the space of finitely additive set functions. To present the main result of [3], let us first introduce the basis notions and notations.

Let (X, d) be a locally compact metric space (with the distance d), $\mathfrak{B}$ be its σ -algebra of Borel sets, $\mathcal{P}(X)$ denote the space of Borel probability measures on X with the weak*-topology, $\mathcal{P}_a(X)$ stand for the set of Borel real-valued normalized finitely additive set functions with the total variation distance (which we will simply call *finitely additive probabilities*), and $C(X)$ the space of continuous maps $\psi : X \rightarrow \mathbb{R}$ with the supremum norm (whose elements are called potentials).

In what follows we will consider a Banach space $\mathbf{B}$ over the field $\mathbb{R}$ equal to either

$$L^\infty(X) = \{\varphi : X \rightarrow \mathbb{R} \mid \varphi \text{ is measurable and bounded}\} \quad (8)$$

or

$$C_c(X) = \{\varphi \in C(X) \mid \varphi \text{ has compact support}\}$$

endowed with the norm $\|\varphi\|_\infty = \sup_{x \in X} |\varphi(x)|$. The Riesz representation theorem asserts that the dual of $C_c(X)$ is identified with the collection of all finite signed measures on $(X, \mathfrak{B})$, whose positive normalized continuous functionals correspond to the space $\mathcal{P}(X)$. It is known that the latter set is compact when equipped with the weak* topology. On the other hand, the dual of $L^\infty(X)$ is isometrically isomorphic to the space of regular finitely additive bounded signed measures on $(X, \mathfrak{B})$ with the total variation norm, whose subset of positive normalized elements is represented by $\mathcal{P}_a(X)$.

Definition 3.1. A function $\Gamma : \mathbf{B} \rightarrow \mathbb{R}$ is called a *pressure function* if it satisfies the following conditions:

$$(C_1) \text{ Increasing: } \varphi \leq \psi \Rightarrow \Gamma(\varphi) \leq \Gamma(\psi) \quad \forall \varphi, \psi \in \mathbf{B}.$$

$$(C_2) \text{ Translation invariant: } \Gamma(\varphi + c) = \Gamma(\varphi) + c \quad \forall \varphi \in \mathbf{B} \quad \forall c \in \mathbb{R}.$$

$$(C_3) \text{ Convex: } \Gamma(t\varphi + (1-t)\psi) \leq t\Gamma(\varphi) + (1-t)\Gamma(\psi) \quad \forall \varphi, \psi \in \mathbf{B}, \forall t \in [0, 1].$$

For a pressure function $\Gamma : \mathbf{B} \rightarrow \mathbb{R}$ we get the following.

$$\begin{aligned} \Gamma(\psi) - \|\varphi - \psi\|_\infty &= \Gamma(\psi - \|\varphi - \psi\|_\infty) \leq \Gamma(\varphi) \leq \Gamma(\psi + \|\varphi - \psi\|_\infty) \\ &= \Gamma(\psi) + \|\varphi - \psi\|_\infty, \end{aligned}$$

$$|\Gamma(\varphi) - \Gamma(\psi)| \leq \|\varphi - \psi\|_\infty \quad \forall \varphi, \psi \in \mathbf{B}.$$

The following is the main abstract variational principle and the upper semi-continuous metric entropy map arising from convex analysis that we will use later.

Theorem 3.2. [3, Theorem 1] Let (X, d) be a metric space and $\Gamma : \mathbf{B} \rightarrow \mathbb{R}$ be a pressure function.

(a) If X is locally compact and $\mathbf{B} = C_c(X)$ then there exists an upper semi-continuous entropy map $\mathfrak{h}$ such that

$$\Gamma(\varphi) = \max_{\mu \in \mathcal{P}(X)} \left\{ \mathfrak{h}(\mu) + \int \varphi d\mu \right\} \quad \forall \varphi \in C_c(X) \quad (9)$$

$$\text{Besides,} \quad \mathfrak{h}(\mu) = \inf_{\varphi \in C_c(X)} \left\{ \Gamma(\varphi) - \int \varphi d\mu \right\} \quad \forall \mu \in \mathcal{P}(X)$$

and if $\alpha : \mathcal{P}_a(X) \rightarrow \mathbb{R} \cup \{+\infty, -\infty\}$ is another function for which (9) holds then $\alpha \leq \mathfrak{h}$. Furthermore, if $\mathcal{E}_\varphi(\Gamma)$ denotes the set of maximizing elements in (9) then there is a residual subset $\mathfrak{R} \subset C_c(X)$ such that $\#\mathcal{E}_\varphi(\Gamma) = 1$ for every $\varphi \in \mathfrak{R}$.

(b) If $\mathbf{B} = L^\infty(X)$, then

$$\Gamma(\varphi) = \max_{\mu \in \mathcal{P}_a(X)} \left\{ \mathfrak{h}(\mu) + \int \varphi d\mu \right\} \quad \forall \varphi \in L^\infty(X) \quad (10)$$

where $\mathfrak{h}(\mu) = \inf_{\varphi \in \mathbf{B}} \left\{ \Gamma(\varphi) - \int \varphi d\mu \right\}$ for every $\mu \in \mathcal{P}_a(X)$. Besides, the map $\mathfrak{h}$ is upper semi-continuous and maximal among any other upper semi-continuous functions attaining the supremum in (10).

4. Topological entropy and topological pressure of NADDS

4.1. Basic definitions and properties

In this section we introduce a topological entropy and topological pressure of NADDS given by a sequence $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ of continuous maps of a compact metric space (X, d) . The sequence of self-maps determines a sequence of metrics

$$d_n(x, y) := \max\{d(f_i \circ f_{i-1} \circ \dots \circ f_1(x), f_i \circ f_{i-1} \circ \dots \circ f_1(y)) : 1 \leq i \leq n-1\},$$

where $x, y \in X$. Following Kolyada and Snoha [13], we define a topological entropy of the NADDS determined by $f_{1,\infty}$ as follows. Fix a positive integer n and $\epsilon > 0$. We say that a subset $A \subset X$ is (n, ϵ) -separated if for any distinct points $a_1, a_2 \in A$

$$d_n(a_1, a_2) := \max\{d(f_i \circ f_{i-1} \circ \dots \circ f_1(a_1), f_i \circ f_{i-1} \circ \dots \circ f_1(a_2)) : 1 \leq i \leq n-1\} \geq \epsilon.$$

Let $s(n, \epsilon, f_{1,\infty}) := \max\{\text{card}(A) : A \text{ is } (n, \epsilon)\text{-separated subset of } X\}.$

Definition 4.1. The quantity

$$h_{top}(f_{1,\infty}) := \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log(s(n, \varepsilon, f_{1,\infty}))$$

is called the *topological entropy* of a NADDs determined by $f_{1,\infty}$. $\square$

Topological pressure of a NADDs, determined by $f_{1,\infty}$, with respect to a potential $\varphi \in C(X)$ we define as follows

$$P_{top}(f_{1,\infty}, \varphi) := \lim_{\varepsilon \rightarrow 0^+} \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_n(f_{1,\infty}, \varphi, \varepsilon),$$

where for any $n \in \mathbb{N}$ and $\varepsilon > 0$

$$P_n(f_{1,\infty}, \varphi, \varepsilon) := \sup_E \left\{ \sum_{x \in E} e^{S_n^{f_{1,\infty}} \varphi(x)} : E \text{ is an } (n, \varepsilon)\text{-separated subset of } X \right\}$$

and

$$S_n^{f_{1,\infty}} \varphi(x) := \varphi(x) + \varphi(f_1(x)) + \varphi(f_2 \circ f_1(x)) + \cdots + \varphi(f_{n-1} \circ f_{n-2} \circ \cdots \circ f_1(x)).$$

Notice that for the potential $\varphi_0 \equiv 0$ one has

$$P_{top}(f_{1,\infty}, \varphi_0) = h_{top}(f_{1,\infty}).$$

From the definition of pressure we get for any $\varphi \in C(X)$ the following inequalities

$$h_{top}(f_{1,\infty}) + \inf_{x \in X} \varphi(x) \leq P_{top}(f_{1,\infty}, \varphi) \leq h_{top}(f_{1,\infty}) + \sup_{x \in X} \varphi(x).$$

The topological pressure has the following properties.

Lemma 4.2. If $\varphi \leq \psi \Rightarrow P_{top}(f_{1,\infty}, \varphi) \leq P_{top}(f_{1,\infty}, \psi)$, for any $\varphi, \psi \in C(X)$.

Proof. Let $n \in \mathbb{N}$ and $x \in X$. If $\varphi \leq \psi$, then

$$\begin{aligned} S_n^{f_{1,\infty}} \varphi(x) &= \varphi(x) + \varphi(f_1(x)) + \varphi(f_2 \circ f_1(x)) + \cdots + \varphi(f_{n-1} \circ f_{n-2} \circ \cdots \circ f_1(x)). \\ &\leq \psi(x) + \psi(f_1(x)) + \psi(f_2 \circ f_1(x)) + \cdots + \psi(f_{n-1} \circ f_{n-2} \circ \cdots \circ f_1(x)). \\ &= S_n^{f_{1,\infty}} \psi(x), \end{aligned}$$

therefore

$$\begin{aligned} P_n(f_{1,\infty}, \varphi, \varepsilon) &= \sup_E \left\{ \sum_{x \in E} e^{S_n^{f_{1,\infty}} \varphi(x)} : E \text{ is an } (n, \varepsilon)\text{-separated subset of } X \right\} \\ &\leq \sup_E \left\{ \sum_{x \in E} e^{S_n^{f_{1,\infty}} \psi(x)} : E \text{ is an } (n, \varepsilon)\text{-separated subset of } X \right\} \\ &= P_n(f_{1,\infty}, \psi, \varepsilon). \end{aligned}$$

Passing to the limsup with $n \rightarrow \infty$ and limit with $\varepsilon \rightarrow 0^+$ we get

$$P_{top}(f_{1,\infty}, \varphi) \leq P_{top}(f_{1,\infty}, \psi). \quad \square$$

Lemma 4.3. $P_{top}(f_{1,\infty}, \varphi + c) = P_{top}(f_{1,\infty}, \varphi) + c$, for any $\varphi \in C(X)$ and $c \in \mathbb{R}$.

Proof. Notice that

$$\begin{aligned} S_n^{f_{1,\infty}}(\varphi + c)(x) &= \varphi(x) + \varphi(f_1(x)) + \varphi(f_2 \circ f_1(x)) + \cdots + \varphi(f_{n-1} \circ f_{n-2} \circ \cdots \circ f_1(x)) + nc \\ &= S_n^{f_{1,\infty}} \varphi(x) + nc. \end{aligned}$$

Therefore,

$$\begin{aligned} P_n(f_{1,\infty}, \varphi + c, \varepsilon) &= \sup_E \left\{ \sum_{x \in E} e^{S_n^{f_{1,\infty}}(\varphi + c)(x)} : E \text{ is an } (n, \varepsilon)\text{-separated subset of } X \right\} \\ &= e^{nc} \sup_E \left\{ \sum_{x \in E} e^{S_n^{f_{1,\infty}} \varphi(x)} : E \text{ is an } (n, \varepsilon)\text{-separated subset of } X \right\} \\ &= e^{nc} P_n(f_{1,\infty}, \varphi, \varepsilon). \end{aligned}$$

Taking logarithms of both sides and passing to the limsup with $n \rightarrow \infty$ and limit with $\varepsilon \rightarrow 0^+$ we get

$$P_{top}(f_{1,\infty}, \varphi + c) = P_{top}(f_{1,\infty}, \varphi) + c. \quad \square$$

Lemma 4.4. $P_{top}(f_\infty, t\varphi + (1-t)\psi) \leq tP_{top}(f_\infty, \varphi) + (1-t)P_{top}(f_\infty, \psi)$, $\forall t \in [0, 1]$ and for any $\varphi, \psi \in C(X)$.

Proof. Choose $t \in [0, 1]$, then

$$\begin{aligned} S_n^{f_{1,\infty}}(t\varphi + (1-t)\psi)(x) \\ &= (t\varphi + (1-t)\psi)(x) + \dots + (t\varphi + (1-t)\psi)(f_{n-1} \circ f_{n-2} \circ \dots \circ f_1(x)) \\ &= tS_n^{f_{1,\infty}}\varphi(x) + (1-t)S_n^{f_{1,\infty}}\psi(x). \end{aligned}$$

By the Holder's inequality (with $p = \frac{1}{t}$ and $q = \frac{1}{1-t}$) we get

$$\sum_{x \in E} e^{S_n^{f_{1,\infty}}(t\varphi + (1-t)\psi)(x)} \leq \left(\sum_{x \in E} e^{S_n^{f_{1,\infty}}(\varphi(x))} \right)^t \cdot \left(\sum_{x \in E} e^{S_n^{f_{1,\infty}}(\psi(x))} \right)^{1-t},$$

where E is an (n, ε) -separated subset of X . Therefore

$$P_n(f_{1,\infty}, t\varphi + (1-t)\psi, \varepsilon) \leq P_n(f_\infty, \varphi, \varepsilon)^t \cdot P_n(f_\infty, \psi, \varepsilon)^{1-t}.$$

Taking logarithms of both sides and passing to limsup with $n \rightarrow \infty$ and limit with $\varepsilon \rightarrow 0^+$ we obtain

$$P_{top}(f_{1,\infty}, t\varphi + (1-t)\psi) \leq tP_{top}(f_{1,\infty}, \varphi) + (1-t)P_{top}(f_{1,\infty}, \psi). \quad \square$$

As a consequence we get the following property.

Corollary 4.5. *Given a NADDS determined by $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ with $h_{top}(f_{1,\infty}) < +\infty$, then $P_{top}(f_{1,\infty}, \cdot)$ is a pressure function.*

4.2. Proof of Theorem A

Proof. Since $P_{top}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ is a pressure function the unifying statement of Theorem 3.2 provides an abstract variational principle for a nonautonomous discrete dynamical system determined by $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ and the topological pressure $P_{top}(f_{1,\infty}, \varphi)$, with respect to $\varphi \in C(X)$. $\square$

4.3. Proof of Corollary 2.1

For a pressure function $\Gamma : C(X) \rightarrow \mathbb{R}$ and potential $\phi \in C(X)$ we say that $\mu \in P(X)$ is a *tangent functional* to Γ at ϕ if the inequality $\Gamma(\phi + \psi) - \Gamma(\phi) \geq \int \psi d\mu$ holds for any $\psi \in C(X)$. Let $\mathcal{T}_\phi(\Gamma)$ be the space of tangent functionals to Γ at ϕ and let

$$\mathcal{E}_\phi(\Gamma) := \{\mu \in P(X) : \Gamma(\phi) = \mathfrak{h}(\mu) + \int \psi d\mu\}.$$

Let us recall the definition of the Gateaux differential (or Gateaux derivative), which is a generalization of the concept of directional derivative in differential calculus. Let F be a function on an open set U of a Banach space X into the Banach space Y , the *Gateaux differential* $dF(u; \psi)$ of F at $u \in U$ in the direction $\psi \in X$ is defined as

$$dF(u; \psi) = \lim_{t \rightarrow 0} \frac{F(u + t \cdot \psi) - F(u)}{t}.$$

If the limit exists for all $\psi \in X$, then one says that F is Gateaux differentiable at u .

Now we can provide the proof of Corollary 2.1.

Proof. Fix a nonautonomous discrete dynamical system $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$, with finite topological entropy $h_{top}(f_{1,\infty})$, and a continuous potential $\varphi \in C(X)$. From Theorem A it follows that for the pressure function $P_{top}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ and a continuous potential $\varphi \in C(X)$ there exists a measure $\mu \in P(X)$ such that $P_{top}(f_{1,\infty}, \varphi) = \mathfrak{h}_{f_{1,\infty}}(\mu) + \int_X \varphi d\mu$. It remains to show that this measure μ is uniquely determined. Since the function $P_{top}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ is convex, for any $\phi, \psi \in C(X)$ we obtain that the function

$$t \rightarrow \frac{P_{top}(f_{1,\infty}, \phi + t \cdot \psi) - P_{top}(f_{1,\infty}, \phi)}{t}$$

is increasing and therefore there exist limits:

$$d^+ P_{top}(f_{1,\infty}, \phi)(\psi) := \lim_{t \searrow 0^+} \frac{P_{top}(f_{1,\infty}, \phi + t \cdot \psi) - P_{top}(f_{1,\infty}, \phi)}{t},$$

$$d^- P_{top}(f_{1,\infty}, \phi)(\psi) := \lim_{t \nearrow 0^-} \frac{P_{top}(f_{1,\infty}, \phi + t \cdot \psi) - P_{top}(f_{1,\infty}, \phi)}{t},$$

hence the pressure function $P_{top}(f_{1,\infty}, \cdot)$ is Gateaux differentialble at ϕ .

From Theorem 2 in [3] it follows that for the pressure function $\Gamma(\cdot) = P_{top}(f_{1,\infty}, \cdot)$ we get equality $\mathcal{E}_\phi(\Gamma) = \mathcal{T}_\phi(\Gamma)$, for any $\phi \in C(X)$. Using the Gateaux differentiability of the pressure function $P_{top}(f_{1,\infty}, \cdot)$ and the Corollary 4 in [3] we get that for ϕ there exists a unique tangent functional in $\mathcal{T}_\phi$, that completes the proof. $\square$

Analyzing the proof of Corollary 2.1, it can be seen that for a nonautonomous discrete dynamical system $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$, defined on a compact metric space (X, d) , an abstract pressure function $\Gamma_{f_{1,\infty}} : C(X) \rightarrow \mathbb{R}$ with $\Gamma_{f_{1,\infty}}(0) < \infty$ and a potential $\varphi \in C(X)$ there exists a unique measure μ on X such that $\Gamma_{f_{1,\infty}}(\varphi) = \mathfrak{h}_{f_{1,\infty}}(\mu) + \int_X \varphi d\mu$ if and only if $\Gamma_{f_{1,\infty}} : C(X) \rightarrow \mathbb{R}$ is Gateaux differentiable at φ .

In general case for a nonautonomous dynamical system $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$, there is no common f_n -invariant Borel probability measure, for any $n \in \mathbb{N}$ (see Example 4.6).

Example 4.6. Consider $f_{1,\infty} = \{f_n : S^1 \rightarrow S^1\}_{n=1}^\infty$. Assume that f_i , where $i = 1, 2$, are circle diffeomorphisms, such that each of them has two fixed points: a source p_i and a sink q_i which split S^1 into two arcs A_i and B_i each of them being contracted to q_i . Additionally, we assume that $\{p_1, q_1\} \cap \{p_2, q_2\} = \emptyset$. By Proposition 3.1.6 in [21] any measure μ that is f_i -invariant has to be supported in the set $\{p_i, q_i\}$. Therefore f_1 and f_2 do not have a common invariant measure, which consequently gives that $\mathcal{P}_{f_{1,\infty}}(X) = \emptyset$. $\square$

However, there are many special cases of nonautonomous dynamical systems for which invariant measures exist (see Example 2.2 and Example 2.3).

4.4. Proof of Theorem B

Theorem B generalizes the analogous property established in [3] for a single dynamics. Theorem B result relates the restriction to $\mathcal{P}_{f_{1,\infty}}(X)$ of the variational metric entropy $\mathfrak{h}_{f_{1,\infty}}$ with the star-entropy of NADDS, defined by

$$\mu \in \mathcal{P}_{f_{1,\infty}}(X) \mapsto h_{f_{1,\infty}}^*(\mu) := \sup \left\{ \limsup_{n \rightarrow +\infty} h_{f_{1,\infty}}(\mu_n) \mid \mu_n \in \mathcal{P}_{f_{1,\infty}}(X) \text{ and } \lim_{n \rightarrow +\infty} \mu_n = \mu \right\},$$

where the convergence of $(\mu_n)_{n \in \mathbb{N}}$ takes place in the weak*-topology.

Now we can present a proof of Theorem B.

Proof. Let $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ be a nonautonomous discrete dynamical system, defined on a compact metric space X , and let $P_{top}(f_{1,\infty}, \cdot)$ be the topological pressure function.

(a) As $P_{top}(f_{1,\infty}, \cdot)$ satisfies the variational principle (4) with respect to a common $f_{1,\infty}$ -invariant Borel probability measures μ on X , it is easy to conclude that, for every $\varphi, \psi \in C(X)$ and every f_n , we have

$$P_{top}(f_{1,\infty}, \varphi + \psi \circ f_n - \psi) = P_{top}(f_{1,\infty}, \varphi) = P_{top}(f_{1,\infty}, \varphi + \psi - \psi \circ f_n). \quad (11)$$

Indeed, by the variational principle we get

$$\begin{aligned} P_{top}(f_{1,\infty}, \varphi + \psi \circ f_n - \psi) &= \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int (\varphi + \psi \circ f_n - \psi) d\mu \right\} \\ &= \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} = P_{top}(f_{1,\infty}, \varphi), \end{aligned}$$

where the second equality is due to the assumption that the measure μ is f_n -invariant and similarly with regard to the second equality in (11).

Now we claim that every probability measure attaining the maximum in (2) is $f_{1,\infty}$ -invariant. Consider $\psi \in C(X)$, $f_n : X \rightarrow X$ and fix $\mu_\varphi, \mu_1, \mu_2 \in \mathcal{P}(X)$ provided by (2) such that

$$\begin{aligned} P_{\text{top}}(f_{1,\infty}, \varphi) &= h_{f_{1,\infty}}(\mu_\varphi) + \int \varphi d\mu_\varphi \\ P_{\text{top}}(f_{1,\infty}, \varphi + \psi \circ f_n - \psi) &= h_{f_{1,\infty}}(\mu_1) + \int \varphi d\mu_1 + \int (\psi \circ f_n) d\mu_1 - \int \psi d\mu_1 \\ P_{\text{top}}(f_{1,\infty}, \varphi + \psi - \psi \circ f_n) &= h_{f_{1,\infty}}(\mu_2) + \int \varphi d\mu_2 + \int \psi d\mu_2 - \int (\psi \circ f_n) d\mu_2. \end{aligned}$$

The first two equalities, property (11) and the variational principle (2) now yield

$$\begin{aligned} h_{f_{1,\infty}}(\mu_\varphi) + \int \varphi d\mu_\varphi &= h_{f_{1,\infty}}(\mu_1) + \int \varphi d\mu_1 + \int (\psi \circ f_n) d\mu_1 - \int \psi d\mu_1 \\ &\geq h_{f_{1,\infty}}(\mu_\varphi) + \int \varphi d\mu_\varphi + \int (\psi \circ f_n) d\mu_\varphi - \int \psi d\mu_\varphi \end{aligned}$$

and so $\int (\psi \circ f_n) d\mu_\varphi - \int \psi d\mu_\varphi \leq 0$. In a similar way, we deduce that

$$\begin{aligned} h_{f_{1,\infty}}(\mu_\varphi) + \int \varphi d\mu_\varphi &= h_{f_{1,\infty}}(\mu_2) + \int \varphi d\mu_2 + \int \psi d\mu_2 - \int (\psi \circ f_n) d\mu_2 \\ &\geq h_{f_{1,\infty}}(\mu_\varphi) + \int \varphi d\mu_\varphi + \int \psi d\mu_\varphi - \int (\psi \circ f_n) d\mu_\varphi \end{aligned}$$

so $\int \psi d\mu_\varphi - \int (\psi \circ f_n) d\mu_\varphi \leq 0$. Therefore, for any $n \in \mathbb{N}$ the equality $\int \psi d\mu_\varphi = \int (\psi \circ f_n) d\mu_\varphi$ holds, it means that the measure μ_φ is f_n -invariant, for any $n \in \mathbb{N}$. Thus, μ_φ is $f_{1,\infty}$ -invariant.

(b) The proof of this assertion is similar to the one given in [22, p. 222] for single dynamics. Indeed, from the assumptions that $h_{\text{top}}(f_{1,\infty}) < +\infty$ and that the topological pressure satisfies a variational principle with the non-negative metric entropy map $h_{f_{1,\infty}}$ we conclude the following. We claim that for given $\mu \in \mathcal{P}(X)$,

$$\mu \in \mathcal{P}_{f_{1,\infty}}(X) \Leftrightarrow \int \varphi d\mu \leq P_{\text{top}}(f_{1,\infty}, \varphi) \quad \forall \varphi \in C(X).$$

Indeed, for $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$ the variational principle yields that the inequality $\int \varphi d\mu \leq P_{\text{top}}(f_{1,\infty}, \varphi)$ holds for any $\varphi \in C(X)$.

Now suppose that μ is a finite signed measure such that for any $\varphi \in C(X)$ we have $\int \varphi d\mu \leq P_{\text{top}}(f_{1,\infty}, \varphi)$. In the first step we show that μ takes only non-negative values. To this aim suppose that $\varphi \geq 0$, $\epsilon > 0$ and $n \in \mathbb{N}$. Then

$$\begin{aligned} \int n \cdot (\varphi + \epsilon) d\mu &= - \int -n \cdot (\varphi + \epsilon) d\mu \geq -P_{\text{top}}(f_{1,\infty}, -n \cdot (\varphi + \epsilon)) \\ &\geq -[h_{\text{top}}(f_{1,\infty}) + \sup_{x \in X} (-n \cdot (\varphi + \epsilon))] \\ &= -h_{\text{top}}(f_{1,\infty}) + n \cdot \inf_{x \in X} (\varphi + \epsilon) > 0 \quad \text{for large } n. \end{aligned}$$

Therefore $\int (\varphi + \epsilon) d\mu > 0$. Since ϵ is arbitrary small we get $\int \varphi d\mu \geq 0$ and μ is a measure.

In the next step we show that μ is a probability measure. Take $n \in \mathbb{Z}$, then

$$\int n d\mu \leq P_{\text{top}}(f_{1,\infty}, n) = h_{\text{top}}(f_{1,\infty}) + n.$$

So for $n \in \mathbb{N}$ we get $\mu(X) = \int 1 d\mu \leq 1 + \frac{h_{\text{top}}(f_{1,\infty})}{n}$ and hence $\mu(X) \leq 1$. On the other hand, for negative n we get $\mu(X) = \int 1 d\mu \geq 1 + \frac{h_{\text{top}}(f_{1,\infty})}{n}$ which yields $\mu(X) \geq 1$. Therefore we have proved that $\mu(X) = 1$.

In the next step we show that $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$. For any $m \in \mathbb{Z}$, arbitrary $\varphi \in C(X)$ and fixed $n \in \mathbb{N}$ due to (11) we get

$$m \int (\varphi \circ f_n - \varphi) d\mu \leq P_{\text{top}}(f_{1,\infty}, m \cdot (\varphi \circ f_n - \varphi)) = P_{\text{top}}(f_{1,\infty}, 0) = h_{\text{top}}(f_{1,\infty}).$$

If $m > 1$ then dividing by m and letting m go to ∞ we get $\int (\varphi \circ f_n - \varphi) d\mu \leq 0$. If $m < -1$ then dividing by m and letting m go to $-\infty$ we get $\int (\varphi \circ f_n - \varphi) d\mu \geq 0$.

Therefore for any potential $\varphi \in C(X)$ and arbitrary $n \in \mathbb{N}$ we have $\int \varphi \circ f_n d\mu = \int \varphi d\mu$, so $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$. The claim has been proved.

Therefore, by formula (3) we get

$$\mu \in \mathcal{P}_{f_{1,\infty}}(X) \Leftrightarrow \mathfrak{h}_{f_{1,\infty}}(\mu) \geq 0.$$

Thus we have proved that the values of the map $\mathfrak{h}_{f_{1,\infty}}$ determine the elements of the set $\mathcal{P}_{f_{1,\infty}}(X)$.

(c) By definition, for every $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$ one has $0 \leq h_{f_{1,\infty}}(\mu) \leq h_{f_{1,\infty}}^*(\mu)$. Moreover, by the variational principle (4), for each $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$

$$h_{f_{1,\infty}}(\mu) \leq P_{\text{top}}(f_{1,\infty}, \varphi) - \int \varphi d\mu \quad \forall \varphi \in C(X)$$

and therefore

$$\mathfrak{h}_{f_{1,\infty}}(\mu) = \inf_{\varphi \in C(X)} \left\{ P_{\text{top}}(f_{1,\infty}, \varphi) - \int \varphi d\mu \right\} \geq h_{f_{1,\infty}}(\mu). \quad (12)$$

Consequently, if $h_{f_{1,\infty}}^*(\mu) > \mathfrak{h}_{f_{1,\infty}}(\mu)$ for some $\mu \in \mathcal{P}_{f_{1,\infty}}(X)$, then there would exist $\nu \in \mathcal{P}_{f_{1,\infty}}(X)$ satisfying $h_{f_{1,\infty}}(\nu) > \mathfrak{h}_{f_{1,\infty}}(\nu)$, contradicting (12). This proves that

$$0 \leq h_{f_{1,\infty}}(\mu) \leq h_{f_{1,\infty}}^*(\mu) \leq \mathfrak{h}_{f_{1,\infty}}(\mu) \quad \forall \mu \in \mathcal{P}_{f_{1,\infty}}(X). \quad (13)$$

(d) Observe now that the inequalities (13) imply that

$$\begin{aligned} \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} &\leq \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^*(\mu) + \int \varphi d\mu \right\} \\ &\leq \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\}. \end{aligned}$$

On the other hand, as $\mathcal{P}_{f_{1,\infty}}(X) \subset \mathcal{P}(X)$ and $\mathfrak{h}_{f_{1,\infty}}$ is upper semi-continuous, for every $\varphi \in C(X)$ we have

$$\max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} \leq \max_{\mu \in \mathcal{P}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\}.$$

Therefore, from the variational principles (2) and (4) we obtain

$$\begin{aligned} \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} &\leq \max_{\mu \in \mathcal{P}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} \\ &= P_{\text{top}}(f_{1,\infty}, \varphi) = \sup_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} \\ &\leq \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^*(\mu) + \int \varphi d\mu \right\} \leq \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\}. \end{aligned}$$

Consequently,

$$P_{\text{top}}(f_{1,\infty}, \varphi) = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ \mathfrak{h}_{f_{1,\infty}}(\mu) + \int \varphi d\mu \right\} = \max_{\mu \in \mathcal{P}_{f_{1,\infty}}(X)} \left\{ h_{f_{1,\infty}}^*(\mu) + \int \varphi d\mu \right\}.$$

The proof is complete. $\square$

5. Misiurewicz entropy and Misiurewicz pressure for NADDS

5.1. Basic definitions

In 1976, Misiurewicz [17] defined a topological pressure and a measure-theoretic entropy for a $\mathbb{Z}_+^n$ action on a compact space. His celebrated result says that the variational principle holds for a $\mathbb{Z}_+^n$ action on a compact space. Our definition of Misiurewicz pressure of $f_{1,\infty}$ was inspired by his definition of the topological pressure for a $\mathbb{Z}_+^n$ action [17].

In this section we consider a NADDS $f_{1,\infty} := \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$ defined on a compact space X . For this purpose let $\mathcal{W}$ be the set of all neighborhoods of the diagonal $\{(x, x) : x \in X\} \subset X \times X$ ordered by inclusion. It is known (see [13]) that this is uniform structure for X .

For any $n \in \mathbb{N}$, neighborhood $\delta \in \mathcal{W}$, potential $\phi \in C(X)$ and any finite subset $E \subset X$ we define:

$$\begin{aligned}\delta_n &:= \bigcap_{i=1}^n (f_i \circ \dots \circ f_1 \times f_i \circ \dots \circ f_1)^{-1} \delta, \\ \phi_n(x) &:= \sum_{i=1}^n \phi \circ f_i \circ \dots \circ f_1(x), \\ p(\phi_n, E) &:= \log \sum_{x \in E} \exp \left[\sum_{i=1}^n \phi(f_i \circ \dots \circ f_1(x)) \right].\end{aligned}$$

It is clear that $\delta_n \in \mathcal{W}$ and $\phi_n \in C(X)$. A finite set $E \subset X$ is called:

- (1) $(n, \delta)_M$ -separated if for any distinct points $x, y \in E$ we have $(x, y) \notin \delta_n$,
- (2) $(n, \delta)_M$ -spanning if for any $x \in X$ there exists $y \in E$ such that $(x, y) \in \delta_n$.

Define

$$\begin{aligned}P_{n,\delta}(f_{1,\infty}, \phi) &:= \sup \{p(\phi_n, E) : E \subset X \text{ and } E \text{ is } (n, \delta)_M\text{-separated}\}, \\ P_\delta(f_{1,\infty}, \phi) &:= \limsup_{n \rightarrow \infty} \frac{1}{n} P_{n,\delta}(f_{1,\infty}, \phi).\end{aligned}$$

Definition 5.1. The quantity $P_{Mis}(f_{1,\infty}, \phi) := \sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, \phi)$ is called the *Misiurewicz pressure* of $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n \in \mathbb{N}}$, with respect to a potential $\phi \in C(X)$. Taking the potential $\varphi_0 \equiv 0$ we can define Misiurewicz entropy of $f_{1,\infty}$ as

$$h_{Mis}(f_{1,\infty}) := P_{Mis}(f_{1,\infty}, \varphi_0).$$

5.2. Pressure function

Proposition 5.2. The map $\varphi \mapsto P_{Mis}(f_{1,\infty}, \varphi)$ is a pressure function.

Proof. The proof is based on the three lemmas below. □

Lemma 5.3. If $\varphi, \psi \in C(X)$ and $\varphi \leq \psi$, then $P_{Mis}(f_{1,\infty}, \varphi) \leq P_{Mis}(f_{1,\infty}, \psi)$.

Proof. Choose a neighborhood $\delta \in \mathcal{W}$ and an $(n, \delta)_M$ -separated set E . Then for $x \in E$ and potentials $\varphi, \psi \in C(X)$ with $\varphi \leq \psi$ we get the following inequalities

$$\begin{aligned}\sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] &\leq \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)], \\ \sum_{x \in E} \exp \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] &\leq \sum_{x \in E} \exp \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)], \\ \log \sum_{x \in E} \exp \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] &\leq \log \sum_{x \in E} \exp \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)], \\ \sup_E \left\{ \log \sum_{x \in E} \exp \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] \right\} &\leq \sup_E \left\{ \log \sum_{x \in E} \exp \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)] \right\}.\end{aligned}$$

In other words, we obtained the inequality $P_{n,\delta}(f_{1,\infty}, \varphi) \leq P_{n,\delta}(f_{1,\infty}, \psi)$, which yields

$$P_\delta(f_{1,\infty}, \varphi) = \limsup_{n \rightarrow \infty} \frac{1}{n} P_{n,\delta}(f_{1,\infty}, \varphi) \leq \limsup_{n \rightarrow \infty} \frac{1}{n} P_{n,\delta}(f_{1,\infty}, \psi) = P_\delta(f_{1,\infty}, \psi).$$

Applying the supremum over $\delta \in \mathcal{W}$ to both sides of the inequality we get

$$P_{Mis}(f_{1,\infty}, \varphi) = \sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, \varphi) \leq \sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, \psi) = P_{Mis}(f_{1,\infty}, \psi).$$

The proof of the lemma is complete. □

Lemma 5.4. For any $\varphi \in C(X)$ and $c \in \mathbb{R}$ the following equality holds

$$P_{Mis}(f_{1,\infty}, \varphi + c) = P_{Mis}(f_{1,\infty}, \varphi) + c.$$

Proof. Fix a potential $\varphi \in C(X)$. For a neighborhood $\delta \in \mathcal{W}$, an $(n, \delta)_M$ -separated set E , and $c \in \mathbb{R}$ we get

$$\begin{aligned} \log \sum_{x \in E} \exp \sum_{i=1}^n (\varphi + c)[f_i \circ \dots \circ f_1(x)] &= \log \sum_{x \in E} \exp \{n \cdot c + \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)]\} \\ &= \log \sum_{x \in E} \exp \{n \cdot c\} \cdot \exp \left\{ \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] \right\} \\ &= n \cdot c + \log \sum_{x \in E} \exp \left\{ \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] \right\}. \end{aligned}$$

Therefore

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sum_{x \in E} \exp \sum_{i=1}^n (\varphi + c)[f_i \circ \dots \circ f_1(x)] \\ = c + \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sum_{x \in E} \exp \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] \end{aligned}$$

Applying the supremum over $\delta \in \mathcal{W}$ to both sides of the above equality we get

$$P_{Mis}(f_{1,\infty}, \varphi + c) = P_{Mis}(f_{1,\infty}, \varphi) + c. \quad \square$$

Lemma 5.5. For any $\varphi, \psi \in C(X)$ and $a \in [0, 1]$ we have

$$P_{Mis}(f_{1,\infty}, a\varphi + (1-a)\psi) \leq a \cdot P_{Mis}(f_{1,\infty}, \varphi) + (1-a) \cdot P_{Mis}(f_{1,\infty}, \psi).$$

Proof. Let $\varphi, \psi \in C(X)$ and $a \in [0, 1]$. For a neighborhood $\delta \in \mathcal{W}$, an $(n, \delta)_M$ -separated set $A \subset X$ we get respectively

$$\begin{aligned} \sum_{i=1}^n (a \cdot \varphi + (1-a) \cdot \psi)[f_i \circ \dots \circ f_1(x)] &= a \cdot \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] + (1-a) \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)], \\ \sum_{x \in A} e^{\sum_{i=1}^n (a \cdot \varphi + (1-a) \cdot \psi)[f_i \circ \dots \circ f_1(x)]} &= \sum_{x \in A} e^{a \cdot \sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)] + (1-a) \sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)]} \\ &\leq \left\{ \sum_{x \in A} e^{\sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)]} \right\}^a \cdot \left\{ \sum_{x \in A} e^{\sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)]} \right\}^{1-a}, \end{aligned}$$

where the last inequality is due to Holder's inequality. Thus

$$\begin{aligned} \log \sum_{x \in A} e^{\sum_{i=1}^n (a \cdot \varphi + (1-a) \cdot \psi)[f_i \circ \dots \circ f_1(x)]} \\ \leq a \log \sum_{x \in A} e^{\sum_{i=1}^n \varphi[f_i \circ \dots \circ f_1(x)]} + (1-a) \log \sum_{x \in A} e^{\sum_{i=1}^n \psi[f_i \circ \dots \circ f_1(x)]}. \end{aligned}$$

Taking supremum of both sides over the sets A which are $(n, \delta)_M$ -separated and applying the definition of $P_{n,\delta}(f_{1,\infty}, \phi)$ we obtain

$$P_{n,\delta}(f_{1,\infty}, a \cdot \varphi + (1-a) \cdot \psi) \leq a \cdot P_{n,\delta}(f_{1,\infty}, \varphi) + (1-a) \cdot P_{n,\delta}(f_{1,\infty}, \psi),$$

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} P_{n,\delta}(f_{1,\infty}, a \cdot \varphi + (1-a) \cdot \psi) \\ \leq a \limsup_{n \rightarrow \infty} \frac{1}{n} \cdot P_{n,\delta}(f_{1,\infty}, \varphi) + (1-a) \limsup_{n \rightarrow \infty} \frac{1}{n} P_{n,\delta}(f_{1,\infty}, \psi). \end{aligned}$$

In this way we have obtained the inequalities

$$P_\delta(f_{1,\infty}, a \cdot \varphi + (1-a) \cdot \psi) \leq a \cdot P_\delta(f_{1,\infty}, \varphi) + (1-a) \cdot P_\delta(f_{1,\infty}, \psi),$$

$$\sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, a \cdot \varphi + (1-a) \cdot \psi) \leq a \cdot \sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, \varphi) + (1-a) \cdot \sup_{\delta \in \mathcal{W}} P_\delta(f_{1,\infty}, \psi).$$

The proof of the lemma is complete. $\square$

5.3. Proof of Theorem C

Proof. Since $P_{Mis}(f_{1,\infty}, \cdot) : C(X) \rightarrow \mathbb{R}$ is a pressure function the unifying statement of Theorem 3.2 provides an abstract variational principle for the nonautonomous discrete dynamical system determined by $f_{1,\infty} = \{f_n : X \rightarrow X\}_{n=1}^\infty$ and Misiurewicz pressure $P_{Mis}(f_{1,\infty}, \varphi)$ with respect to $\varphi \in C(X)$. $\square$

5.4. Proof of Corollary 2.4

Proof. The proof of Corollary 2.4 is very similar to the proof of Corollary 2.1. $\square$

5.5. Proof of Theorem D

Theorem D result relates the restriction to $\mathcal{P}_{f_{1,\infty}}(X)$ of the variational metric entropy $h_{f_{1,\infty}}^{Mis}$ with the star-entropy of NADDS, defined by

$$\mu \in \mathcal{P}_{f_{1,\infty}}(X)$$

$$\mapsto h_{f_{1,\infty}}^{*Mis}(\mu) := \sup \left\{ \limsup_{n \rightarrow +\infty} h_{f_{1,\infty}}^{Mis}(\mu_n) \mid \mu_n \in \mathcal{P}_{f_{1,\infty}}(X) \text{ and } \lim_{n \rightarrow +\infty} \mu_n = \mu \right\},$$

where the convergence of $(\mu_n)_{n \in \mathbb{N}}$ takes place in the weak*-topology.

Proof. The proof of Theorem D is very similar to the proof of Theorem B. We leave this proof to the reader as a simple exercise. $\square$

Acknowledgments: The author is grateful to the anonymous referees for reading the manuscript very carefully, for the apposite comments and valuable suggestions that have helped to improve the manuscript.

References

- [1] R. L. Adler, A. G. Konheim, M. H. McAndrew: *Topological entropy*, Trans. Amer. Math. Soc. 114 (1965) 309–319.
- [2] A. Biś: *Topological and measure-theoretical entropies of nonautonomous dynamical systems*, J. Dyn. Diff. Equations 30 (2018) 273–285.
- [3] A. Biś, M. Carvalho, M. Miguel, P. Varandas: *A convex analysis approach to entropy functions, variational principles and equilibrium states*, Comm. Math. Phys. 394/1 (2022) 215–256.
- [4] A. Biś, M. Carvalho, M. Miguel, P. Varandas: *Entropy function for semigroup actions*, Studia Mathematica 285 (2025) 257–288.
- [5] N. N. Bogoliubov, N. M. Krylov: *La théorie générale de la mesure dans son application à l'étude de systèmes dynamiques de la mécanique non-linéaire*, Ann. Math., 2nd Ser. 38/1 (1937) 65–113.
- [6] R. Bowen: *Entropy for group endomorphisms and homogeneous spaces*, Trans. Amer. Math. Soc. 153 (1971) 40–414.
- [7] R. Bowen: *Topological entropy for noncompact sets*, Trans. Amer. Math. Soc. 184 (1973) 125–136.
- [8] M. Brin, A. Katok: *On local entropy*, in: *Geometric Dynamics*, J. Palis et al. (eds.), Lecture Notes in Mathematics 1007, Springer, Berlin (2006) 30–38.
- [9] E. I. Dinaburg: *A connection between various entropy characterizations of dynamical systems (Russian)*, Izv. Akad. Nauk SSSR Ser. Mat. 35 (1971) 324–366.
- [10] D. Feng, W. Huang: *Variational principles for topological entropies of subsets*, J. Funct. Analysis 263 (2012) 2228–2254.

- [11] X. Huang, X. Wen, F. Zeng: *Topological pressure of nonautonomous dynamical systems*, Nonlinear Dyn. Syst. Theory 8 (2008) 43–48.
- [12] Ch. Kawan: *Metric entropy of nonautonomous dynamical systems*, Nonauton. Stoch. Dyn. Systems 1 (2014) 26–52.
- [13] J. L. Kelley: *General Topology*, Van Nostrand, Princeton (1955).
- [14] S. Kolyada, M. Misiurewicz, L. Snoha: *Topological entropy of nonautonomous piecewise monotone dynamical systems on the interval*, Fund. Math. 160 (1990) 161–181.
- [15] S. Kolyada, L. Snoha: *Topological entropy of nonautonomous dynamical systems*, Random Comput. Dyn. 4 (1996) 205–233.
- [16] M. Kong, W. Cheng, B. Li: *Topological pressures for nonautonomous systems*, Chaos Solitons Fractals 76 (2015) 82–92.
- [17] M. Misiurewicz: *A short proof of the variational principle for a $\mathbb{Z}_+^n$ action on a compact space*, in: Proc. Int. Conf. Dynamical Systems in Math. Physics, Rennes 1975, Astérisque 40, Soc. Math. France, Paris (1976) 147–157.
- [18] C. Mouron: *Positive entropy on nonautonomous interval maps and the topology of the inverse limit space*, Topology Appl. 154 (2007) 894–907.
- [19] Ya. Pesin: *Dimension Theory in Dimensional: Contemporary Views and Applications Systems*, The University of Chicago Press, Chicago (1997).
- [20] Ya. Pesin, B. Pitskel: *Topological pressure and the variational principle for noncompact sets*, Funct. Analysis Appl. 18/4 (1984) 307–318.
- [21] P. Walczak: *Dynamics of Foliations, Groups and Pseudogroups*, Birkhäuser, Basel (2004).
- [22] P. Walters: *An Introduction to Ergodic Theory*, Springer, New York (1975).
- [23] L. Xu, X. Zhou: *Variational principles for entropies of nonautonomous dynamical systems*, J. Dyn. Diff. Equations 30/3 (2018) 1053–1062.